

Agricultural fungicide attracts honey bees, study finds

January 8 2018, by Diana Yates



Fungicides are among the top contaminants of honey bee hives and can interfere with the bees' ability to metabolize other pesticides. Credit: L. Brian Stauffer

When given the choice, honey bee foragers prefer to collect sugar syrup laced with the fungicide chlorothalonil over sugar syrup alone,

researchers report in the journal *Scientific Reports*.

The puzzling finding comes on the heels of other studies linking fungicides to declines in [honey](#) bee and wild bee populations. One recent study, for example, found parallels between the use of chlorothalonil and the presence of *Nosema bombi*, a fungal parasite, in [bumble bees](#). Greater chlorothalonil use also was linked to range contractions in four declining bumble bee species.

Other research has shown that European honey bees have a very limited repertoire of detoxifying enzymes and that exposure to one potentially toxic compound - including fungicides - can interfere with their ability to metabolize others.


"People assume that fungicides affect only fungi," said University of Illinois entomology professor and department head May Berenbaum, who led the new research with postdoctoral researcher Ling-Hsiu Liao. "But fungi are much more closely related to animals than they are to plants. And toxins that disrupt physiological processes in fungi can also potentially affect them in animals, including insects."

Some scientists have argued that bees may be less susceptible to agricultural chemicals than laboratory studies suggest because the bees might detect potentially toxic chemicals in the environment and avoid them. But a [2015 study](#) found that European honey bees and at least one species of bumble bee actually prefer food laced with neonicotinoid pesticides.

To test whether foraging honey bees showed a preference for other chemicals they are likely to encounter in the wild, Liao set up two feeding stations in a large enclosure. Foraging honey bees could fly freely from one feeder to the other, choosing to collect either [sugar syrup](#) laced with a test [chemical](#) or sugar syrup mixed with a solvent as

the control. Over the course of the study, she tested honey bee responses to nine naturally occurring chemicals, three fungicides and two herbicides at various concentrations.

The trials revealed that honey bees prefer the naturally occurring chemical quercetin over controls at all concentrations tested.



Entomology professor May Berenbaum, left, and postdoctoral researcher Ling-Hsiu Liao found that honey bees have a slight preference for food laced with the fungicide chlorothalonil at certain concentrations. Credit: L. Brian Stauffer

"That makes sense, because everything the honey bees eat has quercetin in it," Berenbaum said. "There's quercetin in nectar, there's quercetin in

pollen. Quercetin is in honey and beebread, and it's a reliable cue that bees use to recognize food."

To the researchers' surprise, the bees also preferred sugar syrup laced with glyphosate - the active ingredient in Monsanto's Roundup herbicide - at 10 parts per billion, but not at higher concentrations. And while the bees actively avoided syrup containing the fungicide prochloraz, they showed a mild preference for sugar syrup laced with chlorothalonil at 0.5 and 50 parts per billion, but not at 500 ppb.

"The bees are not only not avoiding this fungicide, they're consuming more of it at certain concentrations," Berenbaum said.

Fungicides are among the most prevalent contaminants of honey bee hives, and it is likely the bees themselves are bringing these pesticides into the colony through their food-collecting activities. While perplexing, bees' preferences for some potentially toxic chemicals may be the result of their distinct evolutionary history, Berenbaum said.

"Honey bee foragers are gleaners," she said. "They're active from early spring until late fall, and no single floral source exists for them for that whole season. If they don't have a drive to search out something new, that's going to seriously compromise their ability to find the succession of flowers they need. Unnatural chemicals might be a signal for a new food."

The new findings are worrisome in light of [research](#) showing that exposure to [fungicides](#) interferes with honey bees' ability to metabolize the acaricides used by beekeepers to kill the parasitic varroa mites that infest their hives, the researchers said.

"The dose determines the poison," Berenbaum said. "If your ability to metabolize poisons is compromised, then a therapeutic dose can become

a toxic dose. And that seems to be what happens when honey bees encounter multiple pesticides."

More information: Ling-Hsiu Liao et al, Behavioral responses of honey bees (*Apis mellifera*) to natural and synthetic xenobiotics in food, *Scientific Reports* (2017). [DOI: 10.1038/s41598-017-15066-5](https://doi.org/10.1038/s41598-017-15066-5)

Provided by University of Illinois at Urbana-Champaign

Citation: Agricultural fungicide attracts honey bees, study finds (2018, January 8) retrieved 25 April 2024 from <https://phys.org/news/2018-01-agricultural-fungicide-honey-bees.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.