


# Corralling xenon gas out of waste streams

January 31 2018, by Susan Bauer



Metal organic frameworks, with nanopores and a high affinity for xenon, can

separate the gas from air or waste streams. Xenon has applications in various fields including lighting, space propulsion and medicine. Credit: Pacific Northwest National Laboratory

From space propulsion to lighting to surgical anesthesia, the applications and needs for xenon gas are growing. And the good news is that researchers are advancing the science to more easily remove xenon from waste streams and collect the low amounts of it found in the atmosphere.

Researchers at the Department of Energy's Pacific Northwest National Laboratory are at the forefront of research developing porous nanoscale materials to capture xenon. They report in the journal *Chem* this month, that inexpensive materials called [metal-organic frameworks](#) have been very successful in separating the gas in a way that may make it far less expensive than existing means of producing it.

Currently, industry uses a common but expensive process called cryogenic distillation to separate xenon from other gases or the atmosphere. In that costly process, a lot of energy is used to chill entire gas streams down to far below freezing in order to concentrate the xenon.

"The process we've demonstrated to selectively trap xenon in a MOF can be done at room temperature," said Praveen Thallapally, a materials scientist at PNNL and a corresponding author on the paper. "You pass a mixed gas stream over the MOF materials just one time to capture the xenon and it can be stored long term and easily released for industrial applications when you want to use it."

The paper's authors note that xenon would likely be used more if it was more economical to produce. For instance, they point to reports that

show [xenon](#) is considered a better surgical anesthetic than the existing technology as it is more potent, less risky, more environmentally friendly and potentially recyclable.

Xeon also has applications in lighting, flash lamps, arc lamps, radiation detectors, medical imaging, research imaging with [nuclear magnetic resonance](#), semiconductors, lasers, space propulsion, the search for dark matter and nuclear processing.

MOFs, while nano-sized, have a [high surface area](#) and are full of pores that can suck up gases like sponges suck up water. There are thousands of MOFs that exist and can be created but each need to be tuned or optimized to attract and hold different gases of interest.

**More information:** Debasis Banerjee et al. Xenon Gas Separation and Storage Using Metal-Organic Frameworks, *Chem* (2018). [DOI: 10.1016/j.chempr.2017.12.025](#)

Provided by Pacific Northwest National Laboratory

Citation: Corralling xenon gas out of waste streams (2018, January 31) retrieved 9 April 2024 from <https://phys.org/news/2018-01-corralling-xenon-gas-streams.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--