

Methane hydrate dissociation off Spitsbergen not caused by climate change

January 9 2018



White methane hydrate layers traverse the sediments. Credit: MARUM - Zentrum für Marine Umweltwissenschaften, Universität Bremen; G. Bohrmann

For years, methane emissions from the seabed have been observed in the Arctic Ocean off Spitsbergen. Researchers have proposed that the warming of seawater by climate change is responsible for the release of methane, but this has not been confirmed. Now, an international team reports that post-glacial uplift is the most likely cause of methane hydrate breakdown. The study is published today in the international journal *Nature Communications*.

Methane hydrates, also known as flammable ice, occur in many regions of the oceans. But this product of [methane](#) and water only forms a solid compound under high pressure in cold temperatures. If the pressure is too low or the temperature is too high, the hydrates decompose and the methane is released as gas from the sea floor into the water column. Spitsbergen has been experiencing severe outgassing for several years. Does the methane originate from decomposed [methane hydrates](#)? What is the cause of the dissociation of the hydrates? Warming due to [climate change](#), or other natural processes? An international team of scientists has now been able to answer this question. They have published a report in *Nature Communications*.

"Our investigations show that uplift of the [sea floor](#) in this region caused by the melting of the ice masses since the end of the last ice age is probably the reason for the dissolution of methane hydrate, which is already ongoing for several thousand years," explains Prof. Dr. Klaus Wallmann, first author of the study by GEOMAR Helmholtz Centre for Ocean Research Kiel. "The region has raised more than the sea level has risen, causing a pressure relief, so that the methane hydrates dissociate at the stability limit," Wallmann continues.

For their investigations, the scientists carried out expedition MSM 57 with the German research vessel Maria S. Merian, led by the Center for Marine Environmental Sciences at the University of Bremen. The mobile drilling rig MARUM-MeBo70 was also used for this study.

"With this special device, we were for the first time able to gain long sediment cores in this area," explains chief scientist Prof. Dr. Gerhard Bohrmann from MARUM. "In these cores, we found significant amounts of freshwater that originate from decomposed hydrates," Bohrmann continues. The scientists were able to prove that this process started 8,000 years ago, and therefore cannot be attributed to global warming of the past decades.



The rotary drill bit of the MeBo70 has penetrated a layer of limescale activated in soft sediments. Bright aragon cements typically fill the cavities of the sea carbonates. Credit: MARUM - Zentrum für Marine Umweltwissenschaften, Universität Bremen; G. Bohrmann

In addition to the geochemical analyses, results of a model simulation of ice distribution in the Arctic since the last ice age were used. "The results show that the rate of isostatic uplift at our drill sites after melting exceeded the eustatic sea-level rise throughout the post-glacial period," explains Prof. Bohrmann.

"In other words, the land has risen faster and stronger as the sea level rose, so that the pressure in the [hydrate](#) reservoir decreased and the hydrates finally became unstable," adds Prof. Wallmann. Thus, the scientists argue that the dissociation of hydrates can be explained by this process, especially since the warming of sea water in deep layers of the ocean is still low.

The investigations off Spitsbergen show a [methane release](#) not caused by climate warming. Further research efforts are necessary at other locations to investigate whether this applies also to other areas of the Arctic or even in middle latitudes.

More information: Klaus Wallmann et al, Gas hydrate dissociation off Svalbard induced by isostatic rebound rather than global warming, *Nature Communications* (2018). [DOI: 10.1038/s41467-017-02550-9](https://doi.org/10.1038/s41467-017-02550-9)

Provided by Helmholtz Association of German Research Centres

Citation: Methane hydrate dissociation off Spitsbergen not caused by climate change (2018, January 9) retrieved 19 April 2024 from <https://phys.org/news/2018-01-methane-hydrate-dissociation-spitsbergen-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.