

## **Permafrost in continuous motion**

## January 25 2018, by Peter Rüegg



Nothing stands still for long here: measurement station on the rock glacier at the foot of the Furggwanghorn mountain. Credit: Thomas Buchli / ETH Zurich

ETH researchers have identified a rock glacier in canton Valais that is degrading and moving very quickly. Fortunately, it does not present an immediate threat to people and infrastructure.

Rock <u>glaciers</u>, containing ice and <u>rock</u> debris as alpine <u>permafrost</u>, are changing rapidly and profoundly in response to climate change. They create the risk of mass movements as they creep downhill, subside, spread and sometimes collapse. The question is whether these glaciers



can potentially pose a threat to the surrounding area, including local residents and infrastructure.

To answer this question, researchers from a number of ETH institutes, and from the ETH spin-off Terrasense, have carried out a detailed study of a rock glacier situated at the foot of the Furggwanghorn mountain in the Turtmann Valley, in the canton of Valais. Their study has just been published in the journal *Permafrost and Periglacial Processes*.

"To identify the potential threat, you first need to ascertain the state of the rock glacier, and understand how it behaves," says Thomas Buchli, a former doctoral researcher in the Chair of Geotechnical Engineering at ETH Zurich. "There are still some gaps in our knowledge here."

Although many studies exist on rock glaciers, scientists have generally focused on measuring their surface and internal temperatures, as well as their surface creep rates. "The interaction between the soil mechanics, the ground temperature and the water inside a rock glacier has never been investigated in such detail," Buchli emphasises. Different measurements must be combined to achieve this. Most research to date has been limited to aspects such as temperature or surface movements, and this information has been compared with other rock glaciers. But these comparisons are not always helpful for understanding an individual rock glacier.





The surface of the rock glacier pictured on the left in the canton of Valais is changing rapidly as it moves downhill at a steadily increasing pace. Credit: ETH Zurich

When writing his dissertation, the geotechnical engineer Buchli collaborated with geophysicists, hydrologists and geologists, using an impressive arsenal of measuring and monitoring equipment that was positioned on and near the Furggwanghorn rock glacier.

Among other things, the researchers used radar and laser measuring devices to characterise the surface. With the help of GPS, they also determined the surface creep velocity of the rock glacier. To investigate



the structure of the ground, the temperature and water content, as well as the internal creep rates, the engineers bored several holes up to 30 metres deep into the permafrost. They inserted thermistors to measure the temperature and also novel inclinometers in these boreholes. Some of the measurements were unique: in one borehole, for example, Buchli was able to detect for the first time continuous creep movements by the rock glacier of up to three metres. The researchers gained additional information on the stratification and structure of the rock glacier with geophysical investigations.

By linking the data together, the researchers were able to obtain a uniquely detailed picture and a ground model that helped to explain the rock glacier's behaviour.

It was striking how quickly and powerfully the surface of the rock glacier moved and changed during the investigation period. Scientists already knew that the Furggwanghorn rock glacier is warming and degrading faster than the majority of other rock glaciers. "Even so, I was surprised by the speed," says Buchli. "Tilting, twisting, subsiding –everything on the rock glacier was somehow in constant motion."

However, the rock glacier did not move as one mass, but in sections. It moved downhill over several metres per year in one part of the frontal area, while the movement in its accumulation zone was only a few decimetres and it was entirely static in another part of the front section.





The surface of the rock glacier showing zones with different creep velocities. Credit: Thomas Buchli / ETH Zurich

Various factors can be considered as drivers of these changes: rainfall, warming air temperatures, <u>water content</u> and of course the temperature and composition of the permafrost. "There are many factors controlling a rock glacier. Isolating just one of them is difficult," says Buchli.

"The Furggwanghorn rock glacier is also relatively warm," the geotechnical engineer says. At the beginning of the measurements, the temperatures in the core of the permafrost were slightly below freezing point. Surface water does not necessarily freeze and flows through a system riddled with pores and cracks. If the permafrost was colder, the water would turn to ice. This permafrost would contain a more



continuous ice phase, and hence be stronger and less permeable.

The temperatures rose continuously in the permafrost during the six years of measurements. They are currently barely above zero in large areas of the rock glacier.

"Already during drilling, we realised that the rock glacier had to be relatively warm, because the drill head kept encountering places with free water," Buchli recalls.

Water flowing through the rock glacier should also be responsible for some of the surface deformations, he says. Water transports heat from the outside into the rock glacier. The permafrost thaws, and causes subsidence in the ground above. However, the researchers have not yet fully understood how the water flows through the rock glacier. "The water flow through a rock glacier is very complex and difficult to investigate," Buchli says.

The researchers could not find any immediate danger for the area below – in this case, the Turtmann valley. It is more likely that the rock glacier, or parts of it, will quietly subside, leaving unfrozen debris behind. "These could trigger a debris flow during very heavy rainfall in the worstcase scenario and then pose a very real threat," says Buchli. In the case of the Furggwanghorn rock glacier, however, the researchers have not found any evidence of a development in this direction. Even so, rock glaciers still require constant monitoring.

**More information:** Thomas Buchli et al. Kinematic investigations on the Furggwanghorn Rock Glacier, Switzerland, *Permafrost and Periglacial Processes* (2018). DOI: 10.1002/ppp.1968



## Provided by ETH Zurich

Citation: Permafrost in continuous motion (2018, January 25) retrieved 6 May 2024 from <u>https://phys.org/news/2018-01-permafrost-motion.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.