


How single-cell archaea determine what direction to swim

January 23 2018



Archaea are unicellular lifeforms without a cell nucleus – like the far better researched bacteria. Credit: Sonja-Verena Albers

Dr. Tessa Quax has identified the structure of a central protein used by archaea to determine the direction to swim. Archaea are single-cell life forms without a nucleus. She also studied which molecular mechanisms are involved in the transmission of signals from the archaea's environment to its motility structure. Quax, who is a researcher in the lab of Prof. Dr. Sonja-Verena Albers at the Institute of Biology II of the University of Freiburg, has published her research in the scientific journal *Proceedings of the National Academy of Sciences*.

Archaea represent one of the three domains of life, along with [bacteria](#) and eukaryotes, the latter of which are organisms that have a nucleus as well as organelles. Archaea are the direct ancestors of eukaryotes, but are similar to bacteria in structure and organization. They thus represent a very important link in the evolution theory. Archaea can adapt to extreme living conditions, such as hot sulfur springs or extremely salty lakes, and they can also be found in the oceans and in our intestines.

Microorganisms are dependent on being able to move actively: If their [living conditions](#) worsen, they are able to move to somewhere more favorable. Like bacteria, archaea are able to sense environmental stimuli and to respond by making a directed movement, called chemotaxis. For this purpose, during evolution they have developed a unique motility structure that is unlike that of bacteria and eukaryotes. Quax is investigating how outer stimuli are transferred via the chemotaxis system to the motility structure.

The organelle responsible for movement in bacteria, the flagellum, has been studied in detail for more than 30 years. The flagellum consists of as many as 50 proteins, which are assembled according to a predetermined order. The result is a whip-like structure (hence its Latin name) of protein threads that functions like a propeller. Anchored to the cell wall and equipped with an "engine" at its tip, it can rotate, which is how the bacteria swims. Until recently, it was believed that archaea also

use flagella to move. However, the team in Albers's lab demonstrated that there are significant structural differences in the motility organelles of bacteria and archaea, and that the archaea's organelle should be renamed "archaellum."

Tessa Quax and her collaborators have also demonstrated that the central chemotaxis protein in archaea fits exactly to the archaeal motility [structure](#). This means that the structural makeup of this [protein](#) is different between archaea and bacteria, although it fulfills the same function in both. This also corresponds with the observation that bacteria and archaea have different motility structures. This research thus offers insight into why [archaea](#) are able to successfully colonize new habitats.

More information: Tessa E. F. Quax et al., "Structure and function of the archaeal response regulator CheY," *PNAS* (2018).

www.pnas.org/cgi/doi/10.1073/pnas.1716661115

Provided by Albert Ludwigs University of Freiburg

Citation: How single-cell archaea determine what direction to swim (2018, January 23) retrieved 19 April 2024 from <https://phys.org/news/2018-01-single-cell-archaea.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.