

Thermal imaging can detect how animals are coping with their environment, avoiding the need for capture, according

January 31 2018



Credit: University of Glasgow

Thermal imaging can detect how animals are coping with their environment, avoiding the need for capture, according to new research.

The technique, which could transform how biologists investigate responses of wild <u>animals</u> to environmental changes, was tested on a population of a small songbird – the blue tit – at the University of



Glasgow's Scottish Centre for Ecology and the Natural Environment (SCENE) at Loch Lomond.

The thermal imaging technique means biologists can learn how animals are responding to their surroundings without having to capture and measure them (necessary with current methods). This is because body <u>temperature</u> can change when animals make physiological adjustments to preserve energy or deal with environmental stressors

The study, published in *Scientific Reports*, found that <u>skin temperature</u> around the eye in the blue tit is lower in birds in poorer condition, and in birds with higher levels of stress hormones in their bloodstream.

In challenging circumstances, such as during poor weather, or when food is scarce, animals need to save energy. One way they can achieve this is by reducing heat production. Challenging conditions also trigger a stress response, which is associated with changes in blood flow around the body – blood is diverted to the areas with the greatest need, increasing core temperature and reducing surface temperature. So, both processes lead to a lower surface temperature.

Corresponding author Dr. Paul Jerem, from the University's Institute of Biodiversity, Animal Health and Comparative Medicine (IBAHCM) said: "These findings are important because understanding physiological processes is key to answering the questions of why animals behave the way they do, and how they interact with each other and their environment."

"Changes in the physiological processes we detected using <u>thermal</u> <u>imaging</u> are generally the first response to environmental challenges. So being able to easily measure them in <u>wild animals</u> means we might be able to identify populations at risk before any decline takes place – a primary goal of conservation."



Dr. Dominic McCafferty, Senior Lecturer at IBAHCM, added: "Current methods of investigating physiological state in free-living species still generally mean animals need to be trapped and handled, something which is difficult, potentially invasive, and limits research to those species which can be easily caught. Additionally, natural patterns of behaviour are also interrupted."

"By looking at this promising alternative strategy—studying body temperature – we are using characteristics we can measure without invasive sampling to infer other underlying <u>physiological processes</u>."

More information: Paul Jerem et al. Eye region surface temperature reflects both energy reserves and circulating glucocorticoids in a wild bird, *Scientific Reports* (2018). DOI: 10.1038/s41598-018-20240-4

Provided by University of Glasgow

Citation: Thermal imaging can detect how animals are coping with their environment, avoiding the need for capture, according (2018, January 31) retrieved 10 May 2024 from https://phys.org/news/2018-01-thermal-imaging-animals-coping-environment.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.