

Distant tropical storms have ripple effects on weather close to home

February 21 2018, by Anne Manning



Credit: Colorado State University

The famously intense tropical rainstorms along the Earth's equator occur thousands of miles from the United States. But atmospheric scientists know that, like ripples in a pond, tropical weather creates powerful waves in the atmosphere that travel all the way to North America and have major impacts on weather in the U.S.

These far-flung, interconnected [weather](#) processes are crucial to making

better, longer-term weather predictions than are currently possible. Colorado State University [atmospheric scientists](#), led by professors Libby Barnes and Eric Maloney, are hard at work to address these longer-term forecasting challenges.

In a new paper in *npj Climate and Atmospheric Science*, the CSU researchers describe a breakthrough in making accurate predictions of weather weeks ahead. They've created an empirical [model](#) fed by careful analysis of 37 years of historical weather data. Their model centers on the relationship between two well-known global weather patterns: the Madden-Julian Oscillation and the quasi-biennial [oscillation](#).

According to the study, led by former graduate researcher Bryan Mundhenk, the model, using both these phenomena, allows skillful prediction of the behavior of major rain storms, called [atmospheric rivers](#), three and up to five weeks in advance.

"It's impressive, considering that current state-of-the-art numerical weather models, such as NOAA's Global Forecast System, or the European Centre for Medium-Range Weather Forecasts' operational model, are only skillful up to one to two weeks in advance," says paper co-author Cory Baggett, a postdoctoral researcher in the Barnes and Maloney labs.

The researchers' chief aim is improving forecast capabilities within the tricky no-man's land of "subseasonal to seasonal" timescales: roughly three weeks to three months out. Predictive capabilities that far in advance could save lives and livelihoods, from sounding alarms for floods and mudslides to preparing farmers for long dry seasons. Barnes also leads a federal NOAA task force for improving subseasonal to seasonal forecasting, with the goal of sharpening predictions for hurricanes, heat waves, the polar vortex and more.

Atmospheric rivers aren't rivers at all

Atmospheric rivers aren't actual waterways, but "rivers in the sky," according to researchers. They're intense plumes of water vapor that cause extreme precipitation, plumes so large they resemble rivers in satellite pictures. These "rivers" are responsible for more than half the rainfall in the western U.S.

The Madden-Julian Oscillation is a cluster of rainstorms that moves east along the Equator over 30 to 60 days. The location of the oscillation determines where atmospheric waves will form, and their eventual impact on say, California. In previous work, the researchers have uncovered key stages of the Madden-Julian Oscillation that affect far-off weather, including atmospheric rivers.

Sitting above the Madden-Julian Oscillation is a very predictable wind pattern called the quasi-biennial oscillation. Over two- to three-year periods, the winds shift east, west and back east again, and almost never deviate. This pattern directly affects the Madden-Julian Oscillation, and thus indirectly affects weather all the way to California and beyond.

The QBO matters

The CSU researchers created a model that can accurately predict atmospheric river activity in the western U.S. three weeks from now. Its inputs include the current state of the Madden-Julian Oscillation and the quasi-biennial oscillation. Using information on how atmospheric rivers have previously behaved in response to these oscillations, they found that the quasi-biennial oscillation matters - a lot.

Armed with their model, the researchers want to identify and understand deficiencies in state-of-the-art numerical weather models that prevent

them from predicting weather on these subseasonal time scales.

"It would be worthwhile to develop a good understanding of the physical relationship between the Madden-Julian Oscillation and the quasi-biennial oscillation, and see what can be done to improve models' simulation of this relationship," Mundhenk said.

Another logical extension of their work would be to test how well their model can forecast actual rainfall and wind or other severe weather, such as tornadoes and hail.

More information: Bryan D. Mundhenk et al, Skillful empirical subseasonal prediction of landfalling atmospheric river activity using the Madden–Julian oscillation and quasi-biennial oscillation, *npj Climate and Atmospheric Science* (2017). [DOI: 10.1038/s41612-017-0008-2](https://doi.org/10.1038/s41612-017-0008-2)

Provided by Colorado State University

Citation: Distant tropical storms have ripple effects on weather close to home (2018, February 21) retrieved 1 May 2024 from <https://phys.org/news/2018-02-distant-tropical-storms-ripple-effects.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--