

Emerald ash borer: How cities and towns can prepare for invasion

February 6 2018



By the time the adult emerald ash borer (*Agrilus planipennis*) appears, it may be too late for nearby ash trees, as the pest inflicts the most damage in its larval form as it burrows beneath the bark of the tree. A new guide published in the open-access *Journal of Integrated Pest Management* outlines a set of four options for communities to choose from as they plan for the impact of the emerald ash borer. Credit: Houping Liu, Ph.D., Pennsylvania Department of Conservation and Natural Resources

Since its discovery in the United States in 2002, the emerald ash borer has swiftly become the most destructive non-native forest pest to ever invade the country. As the insect's range continues to grow, despite ongoing efforts to fight it, communities with ash trees in both urban areas and woodlands are left to pick up the pieces.

Despite the seemingly grim outlook, hope is not lost. In Pennsylvania, where [emerald ash borer](#) (*Agrilus planipennis*) has been present since 2007, municipalities have found successful ash-[management](#) plans under guidance of the state's Department of Conservation and Natural Resources, and they offer a model for other regions to follow. In a new guide published in the Entomological Society of America's open-access *Journal of Integrated Pest Management*, Houping Liu, Ph.D., forest entomologist at the Pennsylvania DCNR, outlines a set of four options for communities to choose from as they plan for the impact of the emerald ash borer:

- **No action** costs nothing up front, but a community is most likely to be burdened with the cost of hazard-tree removal and replanting afterward.
- **Selective management** involves moderate costs for the protection of some high-value [ash trees](#) in a community and is complemented by limited tree removal and replacement.
- **Preemptive management** costs the most up front by eliminating risk from emerald ash borer through complete tree removal before they become infested.
- **Aggressive management** costs more over time, with the least negative social and environmental impact from emerald ash borer infestation.

Thus far, emerald ash borer has defied efforts to contain or suppress its population levels over large areas. Since first appearing in six counties in Michigan in 2002, it has expanded its range, now present in 31 states

plus Ontario and Quebec, Canada. "It is therefore imperative for local communities with limited resources to focus on the protection of valuable individual [trees](#) instead of emerald ash borer population management," Liu says.



In ash trees, signs and symptoms of infestation by the emerald ash borer (*Agrilus planipennis*) include crown thinning and dieback, woodpecker damage, bark splits, eggs under bark, serpentine larval galleries, notched leaflets by adult feeding, D-shaped exit holes, and dead trees. High larval density eventually leads to tree death within two to four years after canopy thinning and branch dieback become evident. Credit: Houping Liu, Ph.D., Pennsylvania Department of Conservation and Natural Resources

The major parts of a community ash-management plan under the Pennsylvania template include tree inventory, monitoring for emerald ash borer, management tools, wood utilization and disposal, replanting, and education and outreach, among others. In the *Journal of Integrated Pest Management* article, Liu summarizes 12 case studies from a state forest, a state park, five cities, and five boroughs in Pennsylvania that developed ash management plans as part of a pilot project.

"While the major components of the plan may stay the same in other areas, each community must tailor the plan to fit its own conditions and management goals," Liu says. "For example, a community with a strong forestry department will be better prepared to draft and execute its own management plan; a community that values its ash trees will be more willing to pull its resources together to combat this problem; ash species to be protected could be different in different areas; and rules and regulations for tree removal, chemical control, and biological control could be different as well."

Liu's guide details the major management methods for emerald ash borer as well as reasons why the pest has outpaced control efforts. "What makes emerald ash borer so difficult to manage is the cryptic nature in its biology, the lack of effective control methods over large areas, and the complexity of quarantine enforcement," he says. However, successful efforts have been found on local scales in eliminating outlier infestations and in treating individual trees with chemical insecticides.

In the future, employing wasps that parasitize the emerald ash borer or cultivating ash trees that are naturally resistant to infestation may become viable options, but more work is needed. For now, Liu says, the best option is: "Be prepared. The devastation is real."

More information: Houping Liu, Under Siege: Ash Management in the Wake of the Emerald Ash Borer, *Journal of Integrated Pest*

Management (2017). [DOI: 10.1093/jipm/pmx029](https://doi.org/10.1093/jipm/pmx029)

Provided by Entomological Society of America

Citation: Emerald ash borer: How cities and towns can prepare for invasion (2018, February 6)
retrieved 19 April 2024 from

<https://phys.org/news/2018-02-emerald-ash-borer-cities-towns.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.