

Infectious disease in hoverflies linked to honeybee health

February 28 2018



Credit: Royal Holloway, University of London

In research published on 28 February, 2018 in *Biology Letters*, scientists from Royal Holloway, University of London, Oxford University and Cornell University have shown for the first time that viruses that are harmful to honeybees are also present in hoverfly pollinators.

Infectious diseases have been identified as a key driver of bee population declines. The new research suggests that hoverflies (Diptera: Syrphidae) are exposed to the same diseases, and may move the infections around when they feed from the same flowers as the honeybees.

Unlike honeybees, hoverflies are very mobile, and can undertake large-scale annual migrations. Thus the study suggests hoverflies have the potential to spread diseases throughout landscapes, or even across entire continents.

Global declines of [insect pollinators](#) jeopardise the delivery of pollination services in both agricultural and natural ecosystems.

It has been well documented in the past the importance of [infectious diseases](#) in bees, but the study shows for the first time the extent to which the diseases are shared with other pollinator groups.

Dr. Emily Bailes, Post Doctoral Research Assistant at Royal Holloway, who led the research, said: "We have seen a decline in bees in the UK for several years, but this study shows for the first time that hoverflies may be moving these diseases much further than bees normally would.

"This is because of their annual migrations across Europe. This could expose local bee populations to new strains of the diseases and make them more likely to become infected, much like different flu strains in humans. We therefore need to think of ways to limit this transfer between species.

"What we don't know yet is whether the hoverflies are just moving the diseases around, or if they are also causing harm to the hoverflies. This is important to find out because hoverflies are also extremely valuable pollinators, and several species are known to be in decline."

During the study, honeybees and four of the most common species of hoverfly were collected from grassland and woodland near Oxford. The samples were screened for the six most common [honeybee](#) viruses, of which three were detected in both honeybees and hoverflies.

Senior author of the paper, Professor Owen Lewis from Oxford University, added: "Most people think of honeybees as the main pollinators of our crops and wildflowers. However, wild insects such as [hoverflies](#) are hugely important too. The new results suggest that the fates of these pollinating insects may be tightly linked, if diseases such as viruses can spread between them."

More information: Emily J. Bailes et al. First detection of bee viruses in hoverfly (syrphid) pollinators, *Biology Letters* (2018). [DOI: 10.1098/rsbl.2018.0001](#)

Provided by Royal Holloway, University of London

Citation: Infectious disease in hoverflies linked to honeybee health (2018, February 28) retrieved 24 April 2024 from <https://phys.org/news/2018-02-infectious-disease-hoverflies-linked-honeybee.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--