

New insight into the molecular weapons of the plant microbiome

February 5 2018




Image depicts Tne2, an NAD-degrading enzyme secreted through the type 6 secretion system of the plant-protective bacterium *Pseudomonas protegens*. Credit: John Whitney, McMaster University

Like all organisms, plants are associated with bacterial communities in which helpful and harmful bacteria compete for dominance. Among the weaponry of these warring bacteria are molecular syringes that some bacteria can use to inject toxins into others. In a study published in the Feb. 2 issue of the *Journal of Biological Chemistry*, researchers at McMaster University in Canada pinpointed the identity of one such toxin used by a soil-dwelling bacterium that protects plants from disease.

The bacterium *Pseudomonas protegens* can kill soil-dwelling plant pathogens, including fungi and bacteria that attack the roots of important crops such as cotton. *Pseudomonas protegens* releases diverse antimicrobial compounds into the soil, but John Whitney was curious specifically about the compounds that it was injecting directly into other bacteria through the type VI secretion system, or T6SS.

"[The T6SS] is this molecular nanomachine that injects [toxic protein](#) into other species of bacteria and kills them," Whitney said. "Plant protective bacteria that have [T6SS] can protect [plants](#) from pathogens better relative to [bacteria] that don't have it."

Jenny Tang and Nathan Bullen, undergraduate students from the University of Waterloo working with Whitney on a co-op work-study assignment, spearheaded the discovery that the toxic protein used by *P. protegens* against other bacteria acts on a molecule found in nearly all living cells: [nicotinamide adenine dinucleotide](#), or NAD⁺.

NAD⁺ is a cofactor, or "helper" molecule, in many biochemical

reactions. By injecting a protein that destroys NAD⁺, *P. protegens* is able to kill other bacteria.

The team then investigated the genome sequences of many other bacteria to see how widespread the strategy of targeting NAD⁺ is in microbial warfare. They found that many bacteria with secretion systems carry genes similar to the one encoding the NAD-targeting [toxin](#).

"We started to see that this isn't just a way of killing that is enacted by plant-protective bacteria," Whitney said. "If you look at the distribution of this (protein) among all sequenced bacteria, it appears that many different bacteria in many different environmental niches use this mode of action to outcompete other bacteria."

The abundance of these toxins in nature raises many questions: How do different bacteria in different environments evolve to resist this toxin? Are NAD-targeting toxins more effective against some bacterial species than others? Understanding the diversity of bacterial weapons is an active area of study among agricultural researchers who would like to develop better ways to fight plant diseases.

"The identification and characterization of antibacterial toxins produced by plant-protective bacteria may one day allow us to engineer these [bacteria](#) to have enhanced ability to suppress pathogens," Whitney said.

More information: Jenny Y. Tang et al, Diverse NADase effector families mediate interbacterial antagonism via the type VI secretion system, *Journal of Biological Chemistry* (2017). [DOI: 10.1074/jbc.RA117.000178](https://doi.org/10.1074/jbc.RA117.000178)

Provided by American Society for Biochemistry and Molecular Biology

Citation: New insight into the molecular weapons of the plant microbiome (2018, February 5)
retrieved 18 April 2024 from

<https://phys.org/news/2018-02-insight-molecular-weapons-microbiome.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.