


Study shines new light on how Salmonella 'die' at low temperatures

February 6 2018



Under mild heat shock, Salmonella dies mainly due to leakage of small cytoplasmic molecules. Credit: Jennifer McCann / Penn State

The most economical way to kill bacteria that cause common food-borne illnesses—mostly caused by *Salmonella enterica*—is heat, but, the mechanisms that kill *Salmonella* at lower temperatures were not fully understood until now, according to a team of researchers.

Bacteria can develop ways to cope with heat shock, so it is important to develop a complete understanding of how heat kills them, the researchers said.

Using droplet-based electrical sensors she developed while a doctoral student at Purdue University, Aida Ebrahimi, assistant professor of electrical engineering, Penn State, determined that mild heat stress at temperatures around 120 degrees Fahrenheit damages the bacteria's [cell wall](#) without rupturing them.

"We had a hypothesis that the *Salmonella* bacterium might die due to leakage of the cell wall," Ebrahimi said. "If you heat them, the lipids that make up the cell wall vibrate. As the cell wall weakens, it can allow small molecules to leak out. Because these small molecules are mostly ionic, we expected a change of the electrical conductance."

In order to prove their hypothesis, the team developed a sensor that was sensitive to the changes in [electrical conductance](#) of the growth medium. As the bacteria's cell wall lost integrity, charged molecules were ejected from the [cells](#) into the solution containing the bacteria, and consequently the electrical conductivity of the solution changed.

The researchers conducted multiple experiments using both wild-type and heat-resistant *Salmonella* bacteria and correlated the electrical results with fluorescence measurement and standard microbiology protocols. The modified bacteria required higher energy to make the cell membranes permeable enough to leak [small molecules](#). The team also studied heating time and heating method, either a slower ramp-up of

heat or a sudden pulse of heat, and found that pulsed heat was more effective at killing bacteria.

The authors of the paper appearing in the current issue of *Biophysical Journal* then developed an analytical model based on their experiments that correlated membrane damage, cytoplasmic leakage and cell death. By better understanding the mechanisms of bacterial death at elevated temperatures, these findings can potentially improve food safety strategies and provide more efficient ways to deactivate bacteria using shorter duration of heating at lower temperatures.

"We know how [high temperatures](#) kill bacteria," Ebrahimi said. "But we wanted to find out why Salmonella died at lower temperatures. There are benefits to using lower temperatures, such as saving energy and retaining better nutritional quality, compared to food heated to high temperatures. But more importantly, [bacteria](#) can develop resistance to heat shock, so it is important to know how they respond to [heat](#) shock."

More information: "Analyzing Thermal Stability of Cell Membrane of Salmonella using Time-Multiplexed Impedance Sensing," *Biophysical Journal* (2018).

Provided by Pennsylvania State University

Citation: Study shines new light on how Salmonella 'die' at low temperatures (2018, February 6) retrieved 20 March 2024 from <https://phys.org/news/2018-02-salmonella-die-temperatures.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
