


Two-stage gas sensor reports on soil dynamics

February 1 2018



A robust two-stage microbial sensor developed at Rice University will help researchers observe gene expression and the bioavailability of nutrients in environments like soil and sediments without disturbing them. Rather than fluoresce, like current biosensors, these release gas to report on their host

microbes' presence and activity. Credit: Ilene Del Valle/Rice University

A gene "genie" developed by Rice University scientists grants researchers valuable data about microbes through puffs of gas from the soil. The latest version is a robust two-stage microbial sensor that will help bioengineers, geobiologists and other researchers observe gene expression and the bioavailability of nutrients in laboratory facsimiles of environments like soil and sediments without disturbing them.

The gas is produced by [microbes](#) genetically engineered to report on both their environment and activity and mixed into soil samples in contained laboratory experiments. One gas that seeps out tells researchers how many of the target microbes are present and a second gas tells researchers what the microbes are doing. Eventually, the Rice team would like the programmed microbes to reveal whether and how they communicate with each other.

Details about the sensors appear in the American Chemical Society journal *ACS Synthetic Biology*.

The ongoing research began in 2015 with a \$1 million grant by the W.M. Keck Foundation and is led by Rice synthetic biologist Jonathan Silberg, biogeochemist Caroline Masiello and graduate student and lead author Hsiao-Ying (Shelly) Cheng. Their goal has been to measure bioactivity in opaque environments, especially those where changing the environment would alter the results.


Silberg said the new gas-emitting microbes operate on the same principle as those that contain two fluorescent proteins; for example, a green-fluorescing protein would tag all the cells in a dish and a red one would light up when triggered by a [microbial activity](#), like protein expression or

proximity of a specific molecule.

"In those systems, you can check the ratio of green to red and know, on average, what the cells are doing," he said. "But that doesn't work in soils."

Currently, researchers measure microbial activity in soil by grinding samples and using processes like high-performance liquid chromatography to quantify their contents. That not only eliminates the opportunity to study the same sample over time, it also limits the scope of the data.

"Our system answers the right question," Masiello said. "Do microbes know these compounds are present, and what are they doing in response to them?"



The two-stage biosensors developed by Rice University graduate student Hsiao-Ying (Shelly) Cheng are programmed to release puffs of gas to report on their presence and activity levels in genetically engineered bacteria like *E. coli*. They mimic the features of fluorescent sensors scientists use to monitor biological activity, but in opaque environments like soil. Credit: Jeff Fitlow/Rice University


In the Rice lab's ratio-metric system, gases that emerge from modified *E. coli* or other microbes could help scientists measure soil development. Ratio-metric means the output of gas is directly proportional to the input, in this case the level of activity the microbe senses.

In one test, *E. coli* was modified to express enzymes that synthesize ethylene and bromomethane. The bacterium continuously made ethylene, which allowed the researchers to monitor the size of the microbe population, but only made bromomethane when triggered by, in this case, the bioavailability of acylhomoserine lactones (AHL), molecules that facilitate signaling between bacteria.

After Cheng put the *E. coli* in agricultural soil and set the temperature to maximize gas signals, she found that adding short- and long-chain AHL did not affect ethylene output but dramatically affected bromomethane. The highest concentration of short-chain AHL increased the bromomethane signal more than an order of magnitude, and long-chain AHL almost two orders of magnitude.

Tests with another bacteria, *Shewanella*, whose native habitat is a sediment, showed similarly robust results. "The dynamic range for sensing chemicals with what Shelly's built is very good," Silberg said. "It

will vary with the organism, but synthetic biology is really about tuning all of that."



Rice University graduate student Hsiao-Ying (Shelly) Cheng led a project to develop two-stage microbial sensors that can observe and report on gene expression and the bioavailability of nutrients in environments like soil and fluids without disturbing them. Credit: Jeff Fitlow/Rice University

"The particularly useful aspect of this work is the potential to distinguish between what's chemically extractable in a marine or soil environment and what a microbe perceives is there," Masiello said. "Just because we can grind up a [soil](#) and measure something doesn't mean that plants or microbes know what's there. These tools are what we need to be able to,

for the first time, measure microbial perception of their environment."

The modified microbes are meant to be used for lab testing and not in the wild. But tests would be much faster than current processes and allow labs to monitor a sample continuously over time. The researchers anticipate applications not only in [synthetic biology](#) and environmental science but also for tracking the environmental fate of gut bacteria being developed for diagnostics and therapeutics.

Going forward, the Rice lab intends to focus its effort on the conditional output part of the sensor. "As we've been building this, people like (Rice bioscientist) Jeff Tabor and others are standardizing the sensing modules," Silberg said. "We're building new output modules that you could then couple to the great diversity of sensors they are building.

"Shelly's really led the way to prove that we can do gas reporting, and she was the first to do it in soils," he said. "She then showed we could do it with horizontal gene transfer as part of our proof of concept, and now this. The tools are just getting there, and I think applications will be next."

More information: Hsiao-Ying Cheng et al. Ratiometric gas reporting: a non-disruptive approach to monitor gene expression in soils, *ACS Synthetic Biology* (2018). [DOI: 10.1021/acssynbio.7b00405](https://doi.org/10.1021/acssynbio.7b00405)

Provided by Rice University

Citation: Two-stage gas sensor reports on soil dynamics (2018, February 1) retrieved 10 April 2024 from <https://phys.org/news/2018-02-two-stage-gas-sensor-soil-dynamics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.