

Double disinfection treatment for safer drinking water

March 7 2018



Double disinfection treatment for safer drinking water. Credit: University of Eastern Finland

In spite of good progress in water hygiene during the recent decades, contaminated water still causes millions of diseases every year. Most of these diseases are caused by enteric viruses, and better water disinfection methods could help prevent discomfort and even save lives. A new study from the University of Eastern Finland shows that a combined disinfection treatment with chlorine and UV radiation can be highly

effective in water disinfection.

The study isolated different coliphages from treated municipal wastewater. Coliphages are viruses that infect E.coli bacteria, and they can be used as models for human enteric viruses in disinfection studies. Nearly half of the isolated coliphages were highly resistant to [chlorine](#) or UV radiation, and this is why neither chlorine nor UV radiation alone were effective against all coliphages.

"This highlights the importance of a combined [treatment](#)," says doctoral student Alyaa Zyara, MSc, from the University of Eastern Finland, who presented the results in her doctoral dissertation.

When coliphages were first exposed to a low chlorine concentration (0.1 or 0.5 mg Cl/L) for 10 minutes followed by low UV radiation (only 22 mWs/cm²), more than 99.9 percent of all of the coliphages studied became inactivated. However, when the order of the treatment was reversed (UV first, chlorine second), disinfection was much less effective.

"It is more effective to first use a low dosage of chlorine followed by a low dosage of UV radiation than to use high chlorine or UV dosages alone. The order of treatment is also important: using UV [radiation](#) first and chlorine second was less effective. In other words, the combination treatment using chlorine first and UV second can be recommended as a disinfection method for viruses."

The study also tested novel UV-LED technology, as UV-LEDs are a new method for disinfecting drinking water. The study used UV-LEDs operating at a wavelength of 270 nm and with a 120 mW irradiation capacity, which haven't been used in [disinfection](#) studies before. As little as 2 minutes of this UV-LED treatment was enough to cause a 90-99.9 percent reduction in the coliphages tested in a 5.2-litre reactor.

Irradiation time of 10 minutes in the same reactor increased the reductions to 99.99—99.999 percent. A traditional mercury UV lamp at a 254 nm wavelength caused similar or slightly higher reductions in 2 or 10 minutes, but the water volume was only 10 millilitres.

"UV-LEDs are a promising method for disinfecting [water](#), since they consume less energy than traditional mercury UV lamps. Furthermore, as UV-LEDs do not contain any mercury, they are safer for the environment."

More information: Alyaa Zyara et al. UV-LEDs Efficiently Inactivate DNA and RNA Coliphages, *Water* (2017). [DOI: 10.3390/w9010046](https://doi.org/10.3390/w9010046)

Provided by University of Eastern Finland

Citation: Double disinfection treatment for safer drinking water (2018, March 7) retrieved 26 April 2024 from <https://phys.org/news/2018-03-disinfection-treatment-safer.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.