

Dogs with noise sensitivity should be routinely assessed for pain by vets

March 20 2018



Credit: CC0 Public Domain

Dogs which show fear or anxiety when faced with loud or sudden noises

should be routinely assessed for pain by veterinarians, a new study has found.

Animal behavioural scientists from the UK and Brazil examined cases of [dogs](#) which had developed a sensitivity to either loudness, different pitches, or sudden noises, and found that those which also had associated musculoskeletal [pain](#) formed a greater sensitivity to [noise](#).

The study suggested that that fear or anxiety about noise could be association between a fear of noises and underlying pain.

The researchers believe that pain, which could be undiagnosed, could be exacerbated when a noise makes the dogs tense up or 'start', putting extra stress on muscles or joints which are already enflamed, causing further pain. That pain is then associated with a loud or startling noise, leading to a sensitivity to noise and avoidance of situations where they had previously had a bad experience—for example a local park, or a louder room in the house.

Researchers say that veterinarians should ensure that all dogs with behaviour problems associated with noise receive a thorough physical examination to see if pain could be a factor in their fear or anxiety, so that undiagnosed pain could be treated, and the behavioural issue tackled. All the dogs that had pain which were treated showed an improvement of their behaviour. This is the first study to explore this phenomenon.

Professor Daniel Mills from the University of Lincoln's School of Life Sciences, said: "Although the average ages of the dogs were similar, the average age of onset of the problem was nearly four years later in the 'clinical cases'. This strong theme of an older age of onset suggests that the pain may develop later in life and that owners seek treatment more readily, perhaps because the appearance of the problem is out of

character in the subject.

"These results are consistent with the suggestion that whenever there is a late age onset to a behaviour problem, medical issues including those related to pain, should be carefully evaluated. It is worth owners being aware that once pain is successfully managed, the previously learned associations with noise may persist and require their own targeted behaviour modification programme."

Researchers assessed two groups of dogs which presented with noise sensitivity: those which had already been diagnosed with underlying [musculoskeletal pain](#) and those which hadn't.

In both cases, the presenting signs of the dogs' behavioural issue included shaking, trembling and hiding, but those with a diagnosed pain issue also showed a higher level of avoidance when it came to places they had a bad experience with noise - for example attempting to avoid a certain area at a park altogether compared with those without pain.

The dogs with the musculoskeletal pain also started to show signs of fear of noises much later in life than the control cases, and were on average four years older than their pain-free counterparts. Noise triggers ranged from fireworks, thunderstorms and aeroplanes, to gunshots, cars and motorbikes.

Veterinary Medicine student Ana Luisa Lopes Fagundes, from Centro Universitário de Belo Horizonte in Brazil, led the research at Lincoln as part of Brazil's Science without Borders scheme.

She said: "The aim of the study was to explore the presenting signs of dogs with generalised noise sensitivity with and without pain in their muscles or joints. We think that dogs with this sort of [chronic pain](#) may experience the noise quite differently, because if the noise makes them

startle it may cause them to tense their muscles and as consequence they feel pain associated with the noise.

"We found that these dogs which had pain do indeed show different signs, in particular they seem to form much wider associations with the noise, for example they would often tend to avoid not just the place where they had the bad experience but much larger areas too. These dogs also tended to avoid other dogs as well. The findings of this study are really important because they contribute to the dog's welfare and improved behaviour as pain could be identified and subsequently treated."

The findings have been published by the journal *Frontiers In Veterinary Science*.

More information: Ana Luisa Lopes Fagundes et al, Noise Sensitivities in Dogs: An Exploration of Signs in Dogs with and without Musculoskeletal Pain Using Qualitative Content Analysis, *Frontiers in Veterinary Science* (2018). [DOI: 10.3389/fvets.2018.00017](https://doi.org/10.3389/fvets.2018.00017)

Provided by University of Lincoln

Citation: Dogs with noise sensitivity should be routinely assessed for pain by vets (2018, March 20) retrieved 27 April 2024 from <https://phys.org/news/2018-03-dogs-noise-sensitivity-routinely-pain.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.