

Laser-heated nanowires produce micro-scale nuclear fusion with record efficiency

March 14 2018, by Anne Manning




The target chamber (front) and ultra-high intensity laser (back) used in the micro-scale fusion experiment at Colorado State University. Credit: Advanced Beam Laboratory/Colorado State University

Nuclear fusion, the process that powers our sun, happens when nuclear reactions between light elements produce heavier ones. It's also happening—at a smaller scale—in a Colorado State University laboratory.

Using a compact but powerful [laser](#) to heat arrays of ordered nanowires, CSU scientists and collaborators have demonstrated micro-scale nuclear [fusion](#) in the lab. They have achieved record-setting efficiency for the generation of neutrons—chargeless sub-atomic particles resulting from the fusion process. Their work is detailed in a paper published in *Nature Communications*, and is led by Jorge Rocca, University Distinguished Professor in electrical and computer engineering and physics. The paper's first author is Alden Curtis, a CSU graduate student.

Laser-driven controlled fusion experiments are typically done at multi-hundred-million-dollar lasers housed in stadium-sized buildings. Such experiments are usually geared toward harnessing fusion for clean energy applications.

In contrast, Rocca's team of students, research scientists and collaborators, work with an ultra fast, high-powered tabletop laser they built from scratch. They use their fast, pulsed laser to irradiate a target of invisible wires and instantly create extremely hot, dense plasmas—with conditions approaching those inside the sun. These plasmas drive fusion reactions, giving off helium and flashes of energetic neutrons.



Top left: A scanning electron microscope image of aligned deuterated polyethylene nanowires. The other panels are 3-D simulations of the nanowires rapidly exploding following irradiation by an ultra-intense laser pulse. Credit: Advanced Beam Laboratory/Colorado State University

In their *Nature Communications* experiment, the team produced a record

number of neutrons per unit of laser energy—about 500 times better than experiments that use conventional flat targets from the same material. Their laser's target was an array of nanowires made out of a material called deuterated polyethylene. The material is similar to the widely used polyethylene plastic, but its common hydrogen atoms are substituted by deuterium, a heavier kind of hydrogen atom.

The efforts were supported by intensive computer simulations conducted at the University of Dusseldorf (Germany), and at CSU.

Making fusion neutrons efficiently, at a small scale, could lead to advances in [neutron](#)-based imaging, and neutron probes to gain insight on the structure and properties of materials. The results also contribute to understanding interactions of ultra-intense laser light with matter.

More information: Alden Curtis et al, Micro-scale fusion in dense relativistic nanowire array plasmas, *Nature Communications* (2018).
[DOI: 10.1038/s41467-018-03445-z](https://doi.org/10.1038/s41467-018-03445-z)

Provided by Colorado State University

Citation: Laser-heated nanowires produce micro-scale nuclear fusion with record efficiency (2018, March 14) retrieved 3 July 2024 from <https://phys.org/news/2018-03-laser-heated-nanowires-micro-scale-nuclear-fusion.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.