

# New pheromone insight may help predict mountain pine beetle outbreaks

March 19 2018

---



Mountain pine beetle attacking a pine tree. Credit: Christine Chiu

Researchers at the University of British Columbia have shed new light

on how mountain pine beetles produce an important pheromone called trans-verbenol, which could aid in efforts to better predict outbreaks.

In recent years, [mountain](#) pine beetles have destroyed more than 25 million hectares of [pine forests](#) in western North America. In a study published today in *Proceedings of the National Academy of Sciences*, scientists have uncovered previously unknown reservoirs of *trans*-verbenol in the bodies of juvenile mountain pine beetles.

"Trans-verbenol is a pheromone that female mountain pine beetles use to attract other insects to a suitable host tree and coordinate large-scale attacks," said Christine Chiu, lead author of the study and graduate student at the Michael Smith Laboratories at UBC. "It was previously assumed that adult females produced *trans*-verbenol by converting toxic compounds found in the resin of new pine [trees](#) they landed on into pheromones. In this study, we found that the beetles have some secrets: they actually accumulate and store *trans*-verbenol during their larval and pupal stages in the brood trees as they grow."

Female beetles burrow into the bark of healthy pine trees to mate and lay eggs, where their developing larvae gradually gut the tree. Chiu analyzed beetle specimens collected in British Columbia at different life stages using gas chromatography.

She found that both male and female [mountain pine beetles](#) accumulate *trans*-verbenol during their development in the brood tree, but only the [adult females](#) retained it until they emerged and dispersed in search of new host trees.

"What we have found is that [female beetles](#) can release *trans*-verbenol independent of whether or not they are in contact with alpha-pinene, the compound found in the resin of new host trees," said Joerg Bohlmann, the principal investigator of the study and professor at the Michael Smith

Laboratories and faculty of forestry. "This shows that there is a certain level of independence on the part of the attacking female bark beetles to produce this compound."

Mountain pine beetles have been spreading into new habitats, migrating east from B.C. into Alberta, and from lodgepole pine to different species like the jack pine.

"Scientists have been making predictions of how well the beetles will do in those new areas based on the chemical profiles of trees in these new areas. The trees may have different ratios or types of alpha-pinene that result in different kinds of *trans*-verbenol, which may have an impact on how attractive it is to the beetles," said Chiu. "Our findings suggest that rather than looking at the chemical profiles of the trees the beetles are attacking, we should perhaps be looking at the brood trees the beetles develop in."

**More information:** Christine C. Chiu et al., "Monoterpenyl esters in juvenile mountain pine beetle and sex-specific release of the aggregation pheromone *trans*-verbenol," *PNAS* (2018).

[www.pnas.org/cgi/doi/10.1073/pnas.1722380115](http://www.pnas.org/cgi/doi/10.1073/pnas.1722380115)

Provided by University of British Columbia

Citation: New pheromone insight may help predict mountain pine beetle outbreaks (2018, March 19) retrieved 9 April 2024 from

<https://phys.org/news/2018-03-pheromone-insight-mountain-beetle-outbreaks.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--