

Ragweed casts shade on soy production

March 28 2018



The experimental plots required weeding to maintain the right ragweed densities, and to remove other weeds. Credit: Amit Jhala

Ragweed, its pollen potent to allergy sufferers, might be more than a source of sneezes. In the Midwest, the plant may pose a threat to soybean production.

Scientists have found that ragweed can drastically reduce [soybean](#) yield.

"It wasn't really a weed we were worried about too much," says Ethann Barnes, a graduate research assistant in agronomy and horticulture at the University of Nebraska-Lincoln. "We didn't expect it to be this competitive."

Weeds compete with crops for light, water, and nutrients. Common ragweed, which is taller than soy, has historically been overlooked as a threat. And little is known about its impact on soy in the Midwest.

So, the scientists struck out to a soybean field near Mead, Nebraska. In 2015 and 2016, they planted soybean and ragweed in late spring. Within the experimental plots, ragweed density ranged from no plants (a weed-free control) to 12 plants per meter (about 39 inches) of the row.

The researchers had two goals: see if ragweed posed a serious threat to soybean, and see if there's a way to estimate the yield loss early in the growing season.

Barnes was surprised by how much the ragweed stifled the soybean in both years. The [soybean crops](#) did worse than in previous studies. One ragweed plant every 1.6 feet of soybean row decreased [soybean yield](#) by 76% in 2015, and by 40% in 2016. And soybean yield was reduced by 95% in 2015 and 80% in 2016 when common ragweed plants were grown only three inches apart in the soybean row.



A ragweed plant rises above neighboring soybean. In the study, ragweed hurt soybean yields by competing for sunlight. Credit: Ethann Barnes

During the experiment, there was plenty of water to go around for both [plants](#). So, the scientists think ragweed mostly hurt soybean by starving it of sunlight.

"Whether I was presenting at conferences, or even just at my thesis defense, everyone was very surprised how big of a deal common ragweed could be," says Barnes.

What's more, it was very hard to predict early in the year how the soybean would fare. Barnes found that not until early August could he

plug ragweed numbers into an equation and accurately predict what the soybean loss would be.

Now, Barnes and his team are sharing this information with growers in the area. "The ultimate goal of this area of science is for growers to count the number of weeds or make a measurement in their field three weeks into the season. From there they could see whether it's financially a viable option to control their weeds or just leave them in the field," says Barnes. By knowing how much damage the weeds might do, farmers can weigh that loss against the cost of killing the weeds.

More studies will be needed to hone in on the dynamics of [ragweed](#)-and other weed—growth. An end goal, he says, is to predict early in the season how weeds will impede crop yields, so farmers can make better decisions on how to manage them. Such estimates could help farmers know if, when, and how much pesticide to apply.

He hopes his study is a step toward that goal. "Hopefully it'll have an immediate impact for farmers, and advance the science of weed competition research."

The study was published in *Agronomy Journal*.

More information: Ethann R. Barnes et al, Common Ragweed (L.) Interference with Soybean in Nebraska, *Agronomy Journal* (2018). [DOI: 10.2134/agronj2017.09.0554](https://doi.org/10.2134/agronj2017.09.0554)

Provided by American Society of Agronomy

Citation: Ragweed casts shade on soy production (2018, March 28) retrieved 2 May 2024 from <https://phys.org/news/2018-03-ragweed-soy-production.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.