


Trappist-1 exoplanets may have too much water to support life


March 20 2018, by Bob Yirka



Modelled χ^2 goodness of fit for the masses of the TRAPPIST-1 planets as a function of the planet's radius and relative H_2O mass fraction in wt% added to the system. Credit: *Nature Astronomy* (2018) doi:10.1038/s41550-018-0411-6


A team of researchers from Arizona State University and Vanderbilt University has found evidence that suggests the exoplanets surrounding the star Trappist-1 may be too wet to support life. In their paper published in the journal *Nature Astronomy*, the group describes using data from prior efforts that focused on determining the mass and diameter of the stars' planets to calculate densities, and from that, used a computer to model the likely building blocks of each.

Last year, scientists discovered the Trappist-1 star system—a red dwarf 39 light years away surrounded by seven [planets](#), all of which are similar in size to Earth. This discovery set off speculation on the possibility of one or more of the planets harboring [life](#). The researchers with this new effort have thrown a wet blanket on such speculation by suggesting that all of the planets have too much water to support life. In modeling the planets, the researchers found that they all have far more water than Earth, from 50 percent of their mass to 10 percent. The Earth, by contrast, is just 0.2 percent water. So much [water](#) likely means there are no exposed land masses, which suggests no geochemical cycles that could promote an atmosphere. Also, a planet covered by very deep oceans would experience extreme mantle pressure preventing rock from moving upward, likely resulting in a runaway snowball effect.




Slice through a model composition of TRAPPIST-1 'f' which contains over 50 percent water by mass. The pressure of the water alone is enough to cause it to become high-pressure ice. The pressure at the water-mantle boundary is so great that no upper mantle is present at all; instead the shallowest rocks would be more like those seen in the Earth's lower mantle. Credit: ASU

The seven planets are classified as rocky, which means they are not gaseous. Also, three of them reside in the "habitable zone," but their star is approximately 2,000 times dimmer than our own, which means that the planets most likely to support life reside very close to their star. But that could be a problem for a couple of reasons—one is that it means the planets are likely tidally locked, resulting in one side always being too hot while the other is too cold. Also, red dwarfs are known to flare a lot, which could spell doom for life on nearby planets.



All seven planets discovered in orbit around the red dwarf star TRAPPIST-1 could easily fit inside the orbit of Mercury, the innermost planet of our solar system. Credit: NASA/JPL- Caltech

The researchers suggest their results could also have implications for theories on how planets develop as they note that all seven of the planets in the Trappist-1 system lie within the "snow line," but the model shows that the [outer planets](#) likely formed beyond that line and migrated inwards over time.



This graph shows the minimum starting distances of the ice-rich TRAPPIST-1 planets (especially f and g) from their star (horizontal axis) as a function of how quickly they formed after their host star was born (vertical axis). The blue line represents a model where water condenses to ice at 170 K, as in our Solar System's planet-forming disk. The red line applies to water condensing to ice at 212 K, appropriate to the TRAPPIST-1 disk. If planets formed quickly, they must have formed farther away (and migrated in a greater distance) to contain significant ice. Because TRAPPIST-1 dims over time, if the planets formed later, they could have formed closer to the host star and still be ice-rich. Credit: ASU

More information: Inward migration of the TRAPPIST-1 planets as inferred from their water-rich compositions, *Nature Astronomy* (2018) [DOI: 10.1038/s41550-018-0411-6](https://doi.org/10.1038/s41550-018-0411-6) , www.nature.com/articles/s41550-018-0411-6

Abstract

Multiple planet systems provide an ideal laboratory for probing exoplanet composition, formation history and potential habitability. For the TRAPPIST-1 planets, the planetary radii are well established from transits, with reasonable mass estimates coming from transit timing variations^{2,3} and dynamical modelling⁴. The low bulk densities of the TRAPPIST-1 planets demand substantial volatile content. Here we show, using mass–radius–composition models, that TRAPPIST-1f and g probably contain substantial (≥ 50 wt%) water/ice, with TRAPPIST-1 b and c being significantly drier (≤ 15 wt%). We propose that this gradient of water mass fractions implies that planets f and g formed outside the primordial snow line whereas b and c formed within it. We find that, compared with planets in our Solar System that also formed within the snow line, TRAPPIST-1b and c contain hundreds more oceans of water.

We demonstrate that the extent and timescale of migration in the TRAPPIST-1 system depends on how rapidly the planets formed and the relative location of the primordial snow line. This work provides a framework for understanding the differences between the protoplanetary disks of our Solar System versus M dwarfs. Our results provide key insights into the volatile budgets, timescales of planet formation and migration history of M dwarf systems, probably the most common type of planetary host in the Galaxy.

[Press release](#)

© 2018 Phys.org

Citation: Trappist-1 exoplanets may have too much water to support life (2018, March 20)
retrieved 20 March 2024 from <https://phys.org/news/2018-03-trappist-exoplanets-life.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--