

Cytoplasmic streaming is involved in the transmission of signals within giant cells in Chara algae

April 16 2018





клетка харовой водоросли



Top right: spatio-temporal distribution of fluorescence during signal propagation.

Top left: the basic equation of the mathematical model describing signal propagation. Credit: Lobachevsky University

Chara algae are ancient plant organisms that are commonly found in freshwater reservoirs and occur, though more rarely, in water bodies with salt water. An unusual feature of this type of algae is the huge size of individual cells, which can reach up to one mm in diameter and up to several centimeters in length. This feature makes Characean algae a unique subject for the study of intracellular signals, which are associated with the appearance of signaling molecules in some parts of the cell (for example, when they are touched or illuminated) and their propagation to other cell areas. As these intracellular signals propagate, they cause various physiological responses, including changes in photosynthesis and respiration.

A paper published in *Functional Plant Biology* focuses on one of the aspects of the intracellular <u>signaling</u> problem—the role of cytoplasm in the transmission of <u>signal molecules</u> inside the Chara algae cells. It is well known that the cytoplasm that fills the living cells is in a state of an almost continuous motion and only stops moving under stress. The researchers advance a hypothesis that such a motion can participate in the transfer of a signal compound within the cell.

To test the proposed hypothesis, a small section of the Chara cell was illuminated with low intensity light, which activated photosynthesis processes and enhanced fluorescence in the illuminated zone. Further



investigation showed that fluorescence intensification was also observed in the dimmed parts of the plant, which were further away from the illuminated zone, in cell regions positioned downstream the cytoplasmic flow.

In contrast, the areas upstream from the illuminated zone did not change their fluorescence. This result shows that signal molecules formed in the illuminated area, even under weak illumination. These molecules propagated with the moving cytoplasm, affecting photosynthesis and enhancing fluorescence in other parts of the alga cell. In the course of additional testing, an electric current was applied to the cell, causing a temporary stoppage of cytoplasmic streaming. Under these conditions, signal propagation significantly slowed down immediately after the action of the electric current; over time, it reached the initial velocity again. The treatment of Chara <u>cells</u> with cytochalasin, a substance that slows the cytoplasmic streaming, resulted in a gradual decrease in the signal propagation velocity.

According to researcher Vladimir Sukhov,, a <u>mathematical model</u> was developed based on the experimental results that describes the propagation of intracellular signal at different velocities of cytoplasm movement. The model also took into account the process of diffusion of the signal compound molecules. "The proposed model describes the experimental results quite well and makes it possible to predict the efficiency of signal transmission under various conditions, including different velocities of cytoplasmic streaming," Sukhov notes.

The results obtained by the team are of fundamental importance, since it was shown for the first time that cytoplasmic streaming can play an important role in the transmission of intracellular signals in <u>plants</u>, and a mathematical model is proposed for a quantitative description of such propagation. At the same time, the results also have significant applications. Since intracellular signals play an important role in adapting



plants to the effects of unfavorable factors, the ability to regulate them provides an additional tool for controlling the resistance of agricultural plants to such factors and, hence, their yield.

The data published in *Functional Plant Biology* show a new potential application for the development of methods to control intracellular signals by regulating the velocity of cytoplasmic movement. It is obvious that the implementation of such methods in practice is quite a challenge. To solve this task, researchers must first make sure that cytoplasmic streaming plays an important role in intracellular signaling, not only in Characean algae, but also in agricultural plants that are of the greatest interest from a practical point of view. Additionally, researchers must find ways to control cytoplasmic streaming in open field or greenhouse conditions. The mathematical model proposed here can become an important tool for solving this problem—the analysis of this model will enable the prediction of the changes in the cytoplasmic streaming parameters that are necessary to obtain the required applied result.

Provided by Lobachevsky University

Citation: Cytoplasmic streaming is involved in the transmission of signals within giant cells in Chara algae (2018, April 16) retrieved 28 June 2024 from https://phys.org/news/2018-04-cytoplasmic-streaming-involved-transmission-giant.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.