

Deforestation in tropical Africa is not as bad as previously thought

April 10 2018, by Julie C. Aleman



Plant remains, preserved in lake sediments like in the Republic of Congo help give accurate deforestation information. Credit: Carla Staver

Deforestation has massive effects on the biosphere. It contributes to [carbon emissions](#), [changes in water cycles](#) and [biodiversity loss](#). The

main cause of deforestation is the conversion of forested lands to agricultural lands.

Tropical forests are home to an exceptional [diversity of flora and fauna](#), and they represent [one of the largest terrestrial carbon stocks](#). This means that tropical deforestation can release a huge amount of carbon into the atmosphere, [contributing to climate change](#). As a result tropical deforestation had been closely monitored for the past decades. The most common way of doing this has been with the use of satellite images.

The results showed that deforestation was raging all over tropical areas during the [1980s](#) but started to reduce in the [1990s and 2000s](#), except in tropical Asia.

Some studies have even suggested that during the past century (since 1900) Africa lost [55% of its tropical forests](#), [Asia 35%](#) and [Amazonia 11%](#).

But these figures weren't calculated from a baseline of [forest](#) extent in 1900. Rather, they were calculated by using the outputs of models fed with current climate data. This means that they are not very accurate.

[In a recent study](#) we show that the loss of forests in tropical Africa in the past century is far less severe than previously estimated. Our analysis shows that forests have, on average, shrunk by 21.7%.

But deforestation has not been homogeneous. Some areas have been hit much worse than others: for example West and East African forests have been reduced by as much as 80% to 90%.

And, in a particularly surprising result, we found that in some areas of Central Africa, forests have in fact encroached onto savannas, resulting in a net forest expansion, although we don't yet know what the causes

are.

[The results of our study](#) represent good news because deforestation rates are not as bad as previously thought. This also has major implications for conservation efforts in both forests and savannas.

Overturning old assumptions

Identifying areas that have been deforested is very important because it can [serve as a basis](#) for carbon mitigation strategies and conservation planning. For example, planting trees helps trap carbon from the atmosphere and can [help fight ongoing climate change](#). But trees should be planted in deforested areas, not in places that [have long been savannas](#).

In Africa huge areas of savannas are being [targeted for planting trees](#) because they were identified as deforested areas. But ecologists – now backed by our research – are [warning that this may not be the case](#).

A new body of literature is developing around the idea that natural disturbances [play a strong role in determining natural vegetation](#), and shows that earlier studies incorrectly labelled ancient savannas as [newly deforested regions](#). These earlier studies suggested that forests were more widespread in Africa in the past, which explains the previous very high deforestation rates.

Our research paints a different picture and shows that these previous analyses have been incorrect.

New spatial modelling

Our conclusions are different because we relied on two sources of

information that allowed us to come up with a much more accurate account:

- paleo-environmental indicators, such as plant remains, preserved in lake sediments or soils, to reconstruct past vegetation; and
- historical accounts and maps.

Using paleo-environmental indicators was a real challenge. This was because paleodata are relatively rare in Africa, especially in West Africa and the Congo Basin and particularly in the Democratic Republic of Congo. This is due to a paucity of sedimentary archives, and sometimes to access issues to study sites because of political instability.

To address this scarcity problem, we also included information from historical maps and documents dating back to around 1900. These sometimes include description of vegetation. For example, we used the map of "[the great forest region](#)" that Henry M. Stanley designed during the relief expedition of Emin Pacha (1886-1889).

By combining these data sources and using a statistical model, we were able to spatially model the forest extent in 1900 and in 2000, and to provide updated deforestation rates. We estimated that forests were not as widespread as previously thought in tropical Africa, resulting in smaller deforestation rates. We found that [tropical forests](#) have [decreased by 21.7%](#), a figure substantially smaller than the 55% previously cited.

In contrast, when looking at local estimates of [deforestation](#) in West and East Africa, our estimates show an almost complete disappearance of forests. We estimated [deforestation rates](#) of 80% and 90% for West Africa and East Africa, respectively.

On the other hand, Central African forests have expanded by 1.4% at the

expense of savannas. Once again, this pattern was not homogeneous: some areas of the Republic of Congo, the Democratic Republic of Congo and Uganda have been deforested, while areas of Central African Republic and Gabon have been afforested.

Implications

This first data-driven reconstruction of historical forest and [savanna](#) distributions [at the continental scale](#) has strong implications for savanna and forest conservation as it challenges some policies currently being pursued.

For example, there is a global effort to increase the number of trees that can trap carbon. But now that an historical baseline exists, it is urgent to focus efforts in areas that have been truly deforested rather than in [areas](#) that have long been savannas. Indeed, planting forest trees in savannas can result in [a decrease in biodiversity and the loss of unique ecosystems](#).

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: Deforestation in tropical Africa is not as bad as previously thought (2018, April 10) retrieved 1 May 2024 from <https://phys.org/news/2018-04-deforestation-tropical-africa-bad-previously.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.