

Extensive seagrass meadows discovered in Indian Ocean through satellite tracking of green turtles

April 11 2018



Green turtle with seagrass in the Caribbean. Credit: Swansea University



Research led by Swansea University's Bioscience department has discovered for the first time extensive deep-water seagrass meadows in the middle of the vast Indian Ocean through satellite tracking the movement of green sea turtles.

A new study by Swansea University and Deakin University academics, published in the recent *Marine Pollution Bulletin*, reported how the monitoring of the turtles—which forage on seagrasses - tracked the species to the Great Chagos Bank, the world's largest contiguous atoll structure in the Western Indian Ocean.

This area lies in the heart of one of the world's largest Marine Protected Areas (MPAs) and the study involved the use of in-situ SCUBA and baited video surveys to investigate the day-time sites occupied by the turtles, resulting in the discovery of extensive monospecific seagrass meadows of *Thalassondendron ciliatum*.

These habitats are critically important for storing huge amounts of carbon in their sediments and for supporting fish populations.

At three sites that extended over 128?km of the Great Chagos Bank, there was a high seagrass cover (average of 74%) at depths to 29 metres.

The mean species richness of fish in the seagrass meadows was 11 species per <u>site</u>, with a mean average of 8-14 species across the aforementioned three sites.





Fish situated on seagrass meadows. Credit: Swansea University

Results showed a high fish abundance as well as a large predatory shark recorded at all sites and given that the Great Chagos Bank extends over approximately 12,500?km and many other large deep submerged banks exist across the world's oceans, the results suggest that deep-water seagrass may be far more abundant than previously suspected.

Reports of seagrass meadows at these depths with high fish diversity, dominated by large top predators, are relatively limited.

Dr Nicole Esteban, a Research Fellow at Swansea University's Biosciences department, said: "Our study demonstrates how tracking marine megafauna can play a useful role to help identify previously unknown seagrass <u>habitat</u>.



"We hope to identify further areas of critical seagrass habitat in the Indian Ocean with forthcoming turtle satellite tracking research."

Dr Richard Unsworth, from Swansea University's Biosciences department, said: "Seagrasses struggle to live in deep waters due to their need for high light, but in these crystal clear waters of Chagos these habitats are booming.

"Given how these habitats are threatened around the world it's great to come across a pristine example of what <u>seagrass meadows</u> should look like."

More information: N. Esteban et al, The discovery of deep-water seagrass meadows in a pristine Indian Ocean wilderness revealed by tracking green turtles, *Marine Pollution Bulletin* (2018). DOI: 10.1016/j.marpolbul.2018.03.018

Provided by Swansea University

Citation: Extensive seagrass meadows discovered in Indian Ocean through satellite tracking of green turtles (2018, April 11) retrieved 27 June 2024 from <u>https://phys.org/news/2018-04-extensive-seagrass-meadows-indian-ocean.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.