

## FAST's first discovery of a millisecond pulsar

April 30 2018







The Gamma-ray sky map and integrated pulse profiles of the new MSP: Upper panel shows the region of the gamma-ray sky where the new MSP is located. Lower panel a) shows the observed radio pulses in a one-hour tracking observation of FAST. Lower panel b) shows the folded pulses from more than nine years of Fermi-LAT gamma-ray data. Credit: Pei Wang/NAOC

China's 500-meter Aperture Spherical radio Telescope (FAST) discovered a radio millisecond pulsar (MSP) coincident with the unassociated gamma-ray source 3FGL J0318.1+0252 in the Fermi Large Area Telescope (LAT) point-source list. This is another milestone of FAST.

FAST, world's largest single-dish radio telescope, operated by the National Astronomical Observatory of the Chinese Academy of Sciences, has discovered more than 20 new pulsars so far. This first MSP discovery was made by FAST on Feb. 27 and later confirmed by the Fermi-LAT team in reprocessing of Fermi data on April 18th.

The newly discovered pulsar, now named PSR J0318+0253, was isolated via the timing of gamma-ray pulsations. This discovery is the first result from the FAST-Fermi LAT collaboration outlined in a MoU signed between the FAST team and Fermi-LAT team.

"This discovery demonstrates the great potential of FAST in pulsar searching, highlighting the vitality of the large aperture radio telescope in the new era," said Kejia Lee, scientist at the Kavli Institute of Astronomy and Astrophysics, Peking University.



Radio follow-up of Fermi-LAT unassociated sources is an effective technique for finding new pulsars. Previous radio observations, including three epochs with Arecibo in June 2013, failed to detect the MSP. In a one-hour tracking observation with the FAST ultra-wide band receiver, the radio pulses toward 3FGL J0318.1+0252 were detected with a spin period of 5.19 milliseconds, an estimated distance of about 4000 light-years, potentially one of the faintest radio MSPs discovered.

Millisecond pulsars are neutron stars that rotate hundreds of times per second. They play an important role in understanding the evolution of neutron stars and the equation of states of condensed matter, and can also be used to detect low-frequency gravitational waves.

The <u>pulsar</u> timing array (PTA) attempts to detect low-frequency gravitational waves from merging supermassive black holes using the long-term timing of a set of stable <u>millisecond pulsars</u>. Pulsar search is the basis of <u>gravitational wave detection</u> through PTAs.

The planned Commensal Radio Astronomy FAST Survey is expected to discover many <u>millisecond</u> pulsars and thus will make significant contribution to the PTA experiment. "The international <u>radio</u>-astronomy community is excited about the amazing FAST telescope, already showing its power in these discoveries. FAST will soon discover a large number of millisecond pulsars and I am looking forward to seeing FAST's contribution to gravitational wave detection," said George Hobbs, scientist of the Commonwealth Scientific and Industrial Research Organization (CSIRO) of Australia and member of the Gravitational Wave International Committee (GWIC).

FAST will be under commissioning until it reaches the designed specifications and becomes a Chinese national facility.



## Provided by Chinese Academy of Sciences

Citation: FAST's first discovery of a millisecond pulsar (2018, April 30) retrieved 26 April 2024 from <u>https://phys.org/news/2018-04-fast-discovery-millisecond-pulsar.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.