

Researchers connect the data to show an accelerating trend for marine heatwaves in our oceans

April 10 2018



The increasing number, intensity and duration of marine heatwaves is likely to have a profound impact on ocean ecosystems and the industries, like fisheries and tourism, that depend on them. Credit: Daniel Hjalmarsson (Unsplash.com)



An international study in *Nature Communications* co-authored by researchers from the ARC Centre of Excellence for Climate Extremes (CLEX) and the Institute of Marine and Antarctic Studies (IMAS) reveals globally marine heatwaves have increased over the past century in number, length and intensity as a direct result of warming oceans.

From 1925-2016, the study found the frequency of marine heatwaves had increased on average by 34% and the length of each heatwave had increased by 17%. Together this led to a 54% increase in the number of marine heatwave days every year.

"Our research also found that from 1982 there was a noticeable acceleration of the trend in marine heatwaves," said lead author, Dr Eric Oliver from Dalhousie University, Canada.

"While some of us may enjoy the warmer waters when we go swimming, these heatwaves have significant impacts on ecosystems, biodiversity, fisheries, tourism and aquaculture. There are often profound economic consequences that go hand in hand with these events."

Some recent examples show just how significant marine heatwave events can be.

In 2011, Western Australia saw a marine heatwave that shifted ecosystems from being dominated by kelp to being dominated by seaweed. That shift remained even after water temperatures returned to normal.

In 2012, a marine heatwave in the Gulf of Maine led to an increase in lobsters but a crash in prices that seriously hurt the industry's profits.

Persistent warm water in the north Pacific from 2014-2016 led to fishery closures, mass strandings of marine mammals and <u>harmful algal</u>



<u>blooms</u> along coastlines. That heatwave even changed large-scale weather patterns in the Pacific Northwest.

More recently still, Tasmania's intense marine <u>heatwave</u> in 2016 led to disease outbreaks and slowing in growth rates across aquaculture industries.

The researchers used a variety of observational datasets to reveal the trend of increasing marine heatwaves, combining satellite data with a range century long datasets taken from ships and various land based measuring stations. They then removed the influences of natural variability caused by the El Nino Southern Oscillation, the Pacific Decadal Oscillation and the Atlantic Multidecadal Oscillation to find the underlying trend.

"There was a clear relationship between the rise in global average seasurface temperatures and the increase in marine heatwaves, much the same as we see increases in extreme heat events related to the increase in global average temperatures," said co-author Prof Neil Holbrook from IMAS at the University of Tasmania.

"With more than 90% of the heat from human caused global warming going into our oceans, it is likely marine heatwaves will continue to increase. The next key stage for our research is to quantify exactly how much they may change.

"The results of these projections are likely to have significant implications for how our environment and economies adapt to this changing world."

More information: Eric C. J. Oliver et al, Longer and more frequent marine heatwaves over the past century, *Nature Communications* (2018). DOI: 10.1038/s41467-018-03732-9



Provided by University of New South Wales

Citation: Researchers connect the data to show an accelerating trend for marine heatwaves in our oceans (2018, April 10) retrieved 30 April 2024 from <u>https://phys.org/news/2018-04-hotter-longer-frequentmarine-heatwaves.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.