

## Twisting laser light offers the chance to probe the nano-scale

April 5 2018



A twisted laser beam hits a nanoscopic U-shaped gold grating which further twists the beam in either a right or left-handed direction. This deflects the beam in many directions and further splits it into its constituent wavelengths across the color spectrum. Credit: Ventsislav Valev



A new method to sensitively measure the structure of molecules has been demonstrated by twisting laser light and aiming it at miniscule gold gratings to separate out wavelengths.

The technique could potentially be used to probe the structure and purity of molecules in pharmaceuticals, agrochemicals, foods and other important products more easily and cheaply than existing methods.

Developed by physicists at the University of Bath, working with colleagues at the University of Cambridge and University College London, the technique relies on the curious fact that many biological and pharmaceutical molecules can be either 'left-handed' or 'right-handed'.

Although such molecules are built from exactly the same elements they can be arranged in <u>mirror images</u> of each other, and this configuration sometimes changes their properties drastically.

Notoriously the morning sickness drug Thalidomide caused birth defects and deaths in babies before it was pulled from the market in the 1960s. Investigation showed that the drug existed in two mirror images - the right-handed form was effective as a morning sickness drug, but the lefthanded form was harmful to foetuses. This is one example of why testing what 'handedness', or chirality, a molecule has is essential for many valuable products.

The research team from the Centre for Photonics and Photonic Materials, and the Centre for Nanoscience and Nanotechnology at the University of Bath, used a special white-light laser built in-house and directed it through several optical components to put a twist on the beam. The twisted laser beam then hits a nano-scopic U-shaped gold grating which serves as a template for the light, further twisting the beam in either a right or left-handed direction. This deflects the beam in many directions and further splits it into its constituent wavelengths across the



colour spectrum.

By carefully measuring the deflected light scientists can detect tiny differences in intensity across the spectrum which inform them about the chirality of the grating the laser beam interacts with.

The study, published in the journal *Advanced Optical Materials*, demonstrates the technique as a proof of principle.

Christian Kuppe, the PhD student who conducted the experiments, said: "At the moment chiral sensing requires high molecular concentrations because you're looking for tiny differences in how the light interacts with the target molecule.

"By using our gold gratings we aim to use a much smaller amount of molecules to conduct a very sensitive test of their handedness. The next step will be to continue to test the technique with a range of well-known chiral <u>molecules</u>.

"We hope that this will become a valuable way to perform really important tests on all sorts of products including pharmaceuticals and other high-value chemicals."

Dr Ventsislav Valev, who oversaw the work, said: "There's a great deal of scientific excitement about miniaturisation and working on nano-sized dimensions at the very small scale. However, in the rush to go as small as possible, some opportunities have been overlooked. Working with chiral nano-gratings is a great example of that."

**More information:** Christian Kuppe et al, Circular Dichroism in Higher-Order Diffraction Beams from Chiral Quasiplanar Nanostructures, *Advanced Optical Materials* (2018). <u>DOI:</u> <u>10.1002/adom.201800098</u>



## Provided by University of Bath

Citation: Twisting laser light offers the chance to probe the nano-scale (2018, April 5) retrieved 3 May 2024 from <u>https://phys.org/news/2018-04-laser-chance-probe-nano-scale.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.