


100-million-year-old liverwort mimicry in insects

April 26 2018



New green lacewing larva from Burmese amber. Credit: Nanjing Institute of Geology and Palaeontology

Camouflage and mimicry are pervasive throughout the biological world


as part of the usual interactions between predators and their prey, allowing both to avoid detection. Among insects, the icons of mimicry include familiar stick and leaf insects, leaf-like moths and katydids.

Liverworts are among the earliest terrestrial plants, and they have been widespread since the Paleozoic. However, [mimicry](#) between insects and liverworts is extremely rare in both modern and fossil ecosystems.

Recently, researchers from China Agricultural University, the Nanjing Institute of Geology and Palaeontology (NIGP) of the Chinese Academy of Sciences and their colleagues reported a new [lacewing](#) species (green lacewing larvae) based on two larvae from Cretaceous Burmese amber (approximately 100 million years old).

These larvae are anatomically modified to mimic coeval liverworts. This discovery represents the first record of liverwort mimicry by fossil insects and brings to light an evolutionary novelty, both in terms of morphological specialization as well as plant-insect interaction. The research was published in *Current Biology*.

These larvae have broadly foliate lateral plates on their thorax and abdomen. It is the only species known among lacewings with distinctive foliate lobes on the larval body. Such morphological modifications grossly match some coeval liverworts. Therefore, the new larvae are the first example of direct mimicry in lacewing larvae.



B, E, G are larvae, the others are livervorts. Credit: Nanjing Institute of Geology and Palaeontology

The morphological specialization in the new chrysopoid larvae is unique and is unknown among any living or fossil lacewings. While the anatomy of these larvae allowed them to avoid detection, the lack of setae or other anatomical elements for entangling debris as camouflage means their sole defense was mimicry, and members of the species could have been stealthy hunters like living and other fossil Chrysopoidea or been ambush predators aided by their disguise.

The present fossils demonstrate a hitherto unknown life-history strategy among these "wolf in sheep's clothing" predators, one that apparently evolved from a camouflaging ancestor but did not persist within the lineage.

Liverworts are a diverse group distributed throughout the world today, including approximately 9,000 extant species. Liverworts have been diverse since the start of the Late Cretaceous, including in the Burmese amber forest, which was a typical wet, tropical rainforest.



Researchers from China and USA reported a new lacewing species (green lacewing larvae) based on two larvae from the Cretaceous Burmese amber (approximately 100 million years old). These larvae are anatomically modified to mimic coeval liverworts. This discovery represents the first record of liverwort mimicry by fossil insects and brings to light an evolutionary novelty, both in terms of morphological specialization as well as plant-insect interaction. Credit: YANG Dinghua

Like their extant counterparts, Cretaceous liverworts grew on the leaves and bark of trees as well as on other plant surfaces. Therefore, the [larvae](#) most probably lived on trees densely covered by liverworts, with their liverwort mimicry aiding their survival.

More information: *Current Biology* (2018). [DOI: 10.1016/j.cub.2018.03.060](#)

Provided by Chinese Academy of Sciences

Citation: 100-million-year-old liverwort mimicry in insects (2018, April 26) retrieved 2 May 2024 from <https://phys.org/news/2018-04-million-year-old-liverwort-mimicry-insects.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--