

Mississippi River diversions will produce new land, but slowly, study says

April 11 2018



Louisiana 1, a state highway, abruptly ends in open water at the now-coastal community of Leeville, Louisiana in the Mississippi Delta. A new highway (overhead) has been constructed to connect a major industrial port (Port Fourchon) and communities below Leeville with the mainland. Credit: Elizabeth Chamberlain

Although river diversions that bring land building sediment to shrinking

coastlands are the best solution to sustaining portions of the Mississippi Delta, a new Tulane University study concludes that the rate of land building will likely be dwarfed by the rate of wetland loss.

The study, published in the open-access journal *Science Advances*, used optical dating to measure how fast the delta shoreline migrated seaward under natural conditions.

"Optical dating determines when sediment grains were deposited by measuring their last exposure to daylight," said Elizabeth Chamberlain, lead author and recent Tulane PhD graduate. "This method allowed us to date the shoreline of the Lafourche lobe in the central Mississippi Delta and to calculate that it advanced at a rate of 300-500 feet per year for almost 1,000 years."

Prior to human influence, the Mississippi Delta grew at a rate of 2 to 3 square miles per year. But these rates fall far short of the rates of land loss in coastal Louisiana, which have averaged 15 to 20 square miles per year over the past century, the researchers said.

Tulane Vokes Geology Professor Torbjörn Törnqvist, a co-author of the study, said that given the accelerating rates of sea-level rise that will likely continue long into the future, even the best designed river diversions won't be able to prevent more land loss.



Zhixiong Shen (left) and Elizabeth Chamberlain (right) collecting a sample for optical dating along Bayou Lafourche in Louisiana, an abandoned course of the Mississippi River. Credit: Anjali Fernandes

"Therefore difficult choices will have to be made about where to locate these diversions," he said.

With limited federal and state funding, positioning river diversions in areas that have the greatest land building potential and protect the largest population centers, as opposed to multiple diversions targeting the entire shoreline, is the only viable option.

Chamberlain, now a postdoctoral fellow at Vanderbilt University, and Törnqvist conducted the research with colleagues from Coastal Carolina University, the University of Liverpool in the United Kingdom and Wageningen University in The Netherlands.

Justin Lawrence, program director in the National Science Foundation's Division of Earth Sciences, which funded the research, said the results of the study are striking. "The findings will inform coastal policy and management in this low-lying U.S. region."

More information: E.L. Chamberlain et al., "Anatomy of Mississippi Delta growth and its implications for coastal restoration," *Science Advances* (2018). DOI: 10.1126/sciadv.aar4740 , advances.sciencemag.org/content/4/4/eaar4740

Provided by Tulane University

Citation: Mississippi River diversions will produce new land, but slowly, study says (2018, April 11) retrieved 26 April 2024 from <https://phys.org/news/2018-04-mississippi-river-diversions-slowly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.