


# Peptide induces chirality evolution in a single gold nanoparticle

April 19 2018

---



Schematic geometry of chiral gold nanoparticle, mirror imaged twins, synthesized with the chiral peptide additives. Credit: Ki Tae Nam Research Group, Seoul National University


For the first time, scientists have successfully created optically active, chiral gold nanoparticles using amino acids and peptides. Many chemicals significant to life have mirror-image twins (left-handed and right-handed structures), a characteristic that is conventionally called chirality. This study describes how chirality, which is typically observed in organic molecules, can be extended to three-dimensional metallic nanostructures. This newly discovered synthesis method was described in *Nature* and was featured on the cover.

The Korean research teams at Seoul National University (SNU), Pohang University of Science and Technology (POSTECH) and LG Display (LGD) demonstrated the direct transfer of peptide handedness to nanoparticle morphology. Mirror-image peptide twins induced the opposite twist of the chiral nanoparticles, which are further tunable with sequence variation. The chiral gold nanoparticles with different handedness interacted differently with circularly polarized visible light, displaying extensive color modulation. As a result, color change is possible by controlling the light polarization, which has potential applications in future displays.

In newly synthesized gold [nanoparticles](#), chiral elements are arranged on cube-like structures with a side length of only about 100 nm. They can be easily dispersed in solutions and deposited on the substrates while maintaining high chiro-optical activities.

"Based on our understanding of the interface between [peptides](#) and inorganic materials, we have built a new platform technology to control the crystallographic asymmetry," explains Professor Ki Tae Nam at SNU, who led this collaborative project. He added, "This finding can make a direct and immediate impact on optical devices and could be further applied for the development of enantioselective bioinspired catalysts in the near future."

"The potential applications include active color displays, holography, chirality sensors and all-angle negative refractive index materials," said Professor Junsuk Rho at POSTECH, the co-corresponding author.



Electron microscope image of synthesized chiral gold nanoparticles (LEFT), where chiral components are highlighted in blue. Potential application in a color display as a three dimensional polarizer. (Right) Credit: Ki Tae Nam Research Group, Seoul National University

**More information:** Hye-Eun Lee et al, Amino-acid- and peptide-directed synthesis of chiral plasmonic gold nanoparticles, *Nature* (2018).  
[DOI: 10.1038/s41586-018-0034-1](https://doi.org/10.1038/s41586-018-0034-1)

Provided by Seoul National University

Citation: Peptide induces chirality evolution in a single gold nanoparticle (2018, April 19)  
retrieved 2 May 2024 from

<https://phys.org/news/2018-04-peptide-chirality-evolution-gold-nanoparticle.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--