

Conservation through religion? Scientists confirm that sacred natural sites confer biodiversity advantage

April 23 2018



'Our Lady of the Nightingales' - the sacred forest of Panagia Aidonolaloussa. Branches, roots and holes of giant old oaks, Judas-trees, mahaleb cherries and hornbeams bring a Tolkienesque air to this ancient woodland. Credit: K. Stara

Sacred natural sites (SNS) are found all over the world. They are thought to play an important role in conservation but until recently there was little systematic investigation of this claim. Now, new research published in the journal *Biological Conservation* by an international and multidisciplinary team, led by the University of Ioannina and including Bangor University, has shown that there is a notable conservation benefit to SNS. The researchers of the project, known as THALIS-SAGE, chose for their study the region of Epirus, in north-western Greece, that is host to numerous sacred groves protected through religion for hundreds of years.



By studying a wide range of plant and animal taxonomic groups in eight sacred natural sites and comparing their diversity with nearby control sites, mostly reforested areas, the researchers found that the SNS have a small but persistent biodiversity advantage. This advantage is expressed in a number of ways, the most clear indication being the existence of more distinct communities of species amongst the sacred groves than amongst control sites (a measure called beta diversity).

The most notable species group for having higher biodiversity in the sacred natural sites than the control sites was the fungi, often growing in the dead wood of old trees. For the passerine birds (perching birds, which include many song birds) that have been designated as having conservation importance at a European level, twice as many species (eight) were present in the sacred natural sites as in the control sites (four).

It is often asserted that the conservation benefits of SNS are marginal owing to their usually-small sizes. However, this study finds the influence of size to be relatively weak and shows that even small SNS can play a considerable role in biodiversity conservation.

Despite their role in conserving biodiversity, the SNS of Epirus are now themselves endangered. This is due to changing demographics and land use, primarily through rural depopulation in most mountainous areas of Epirus since WW2. Given recent rural depopulation, it is difficult for local communities to continue to enforce the rules, which used to protect their SNS as locally adapted management systems. For this reason, the THALIS-SAGE project also included initiatives for the promotion of their cultural and educational value. Bangor University project partner, Professor John Healey, of the School of Environment, Natural Resources & Geography and University of Ioannina project leader, Professor John Halley, conclude that "the best conservation strategy for these and other SNS of similar sizes worldwide is to link them in networks within



conventional <u>conservation</u> schemes." Project researcher, Dr. Kalliopi Stara, added "this needs to be done in collaboration with local communities."

More information: D.N. Avtzis et al. Quantifying the conservation value of Sacred Natural Sites, *Biological Conservation* (2018). DOI: 10.1016/j.biocon.2018.03.035

Provided by Bangor University

Citation: Conservation through religion? Scientists confirm that sacred natural sites confer biodiversity advantage (2018, April 23) retrieved 17 July 2024 from <u>https://phys.org/news/2018-04-religion-scientists-sacred-natural-sites.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.