

U.S. rivers are becoming saltier – and it's not just from treating roads in winter

April 17 2018, by Sujay Kaushal, Gene E. Likens, Michael Pace And Ryan Utz



Little Missouri River, North Dakota. Credit: Justin Meissen, CC BY-SA

The United States has made enormous progress in reducing water pollution since the Clean Water Act was passed nearly 50 years ago.

Rivers no longer [catch fire](#) when oil slicks on their surfaces ignite. And many harbors that once were [fouled with sewage](#) now draw swimmers and boaters.

But as Earth Day approaches, it is important to realize that new, more complex challenges are emerging. In a [study](#) published earlier this year, we found that a cocktail of chemicals from many human activities is making U.S. rivers saltier and more alkaline across the nation. Surprisingly, [road salt](#) in winter is not the only source: construction, agriculture, and many other activities also play roles across regions.


These changes pose serious threats to [drinking water](#) supplies, urban infrastructure and natural ecosystems. Salt pollution is not currently regulated at the federal level, and state and local controls are inconsistent.

Our research shows that when salts from different sources mix, they can have broader impacts than they would individually. It also shows the importance of supporting water quality monitoring nationwide, so that we can detect and address other pollution problems that have yet to be recognized.

Altered waters

Our group has been studying freshwater salinization for over 15 years. In 2005 we published a [paper](#) that demonstrated that levels of sodium chloride (common table salt) were rapidly increasing in fresh waters across the northeastern United States.

Where is Earth's Water?



Only about 3 percent of the Earth's water supply is fresh water, and only a fraction of that amount is available as liquid water. Credit: USGS

Until that time, scientists thought that salinization was a serious problem mainly in arid regions where water evaporates rapidly, leaving salts behind. But we found that it was affecting major drinking water supplies, exceeding toxic levels for some aquatic organisms and persisting in the environment year-round, even in humid regions.


The main cause we found was the spread of paved surfaces, such as

roads and parking lots. Communities in cold regions use de-icing salts to clear snow from roads during winter, and the more roads they build, the more treatment is needed. We found that a 1 percent increase in paved surfaces could boost salt concentrations in nearby water bodies to levels more than 10 times higher than pristine forested conditions.

In 2013, we published another [study](#) showing that rivers were becoming more alkaline across regions of the eastern United States. At that time [acid rain](#) – i.e., too much acid in rainwater, caused by air pollution – had been a well-known environmental issue for several decades. However, alkalization was not recognized in the same way, and its effects are still poorly understood now.

Alkalinization is the opposite of acidification: It occurs when water's pH value increases instead of falling. As water becomes more alkaline, certain chemicals dissolved in it can become toxic. For example, ammonium is a nutrient in freshwater ecosystems, but is [converted to toxic ammonia gas](#) in significant concentrations in waters with a high pH. Alkaline conditions also enhance release of phosphorus from sediments, which can trigger nuisance blooms of algae and bacteria.

We found that a process we called "[human-accelerated weathering](#)" was breaking down rock and releasing minerals into rivers that were making them more alkaline. The process of weathering rocks and minerals that become exported to rivers is typically slow, but we showed that land development and decades of exposure to acid rain were speeding it. We also suggested that widespread use of geologic materials in fertilizers and concrete was a factor.



Some pH values of common substances. Credit: USEPA

Identifying freshwater salinization syndrome


Our study on human-accelerated weathering showed that along with sodium chloride, other dissolved salts were increasing in [fresh water](#) across large regions of the eastern United States. This made us wonder whether there could be a link to our previous work on salinization in these regions.

We started to recognize that in theory, salt pollution and human-accelerated weathering could be sending increasing quantities of salts that were alkaline into rivers throughout the nation, and that this could increase their pH levels. We knew that ocean water, which is naturally salty, has a higher pH than fresh water because it has accumulated high levels of alkaline salts. After much analysis, we proposed that similar interconnected processes could influence salinity and pH in fresh water.


Many sources release alkaline salts into the environment, including weathering of impervious surfaces, fertilizer and lime use in agriculture,

mine drainage, irrigation runoff and winter use of road salt. Initially, parts of these alkaline salts bind to soil. But when they come into contact with sodium – for example, excess road salt – chemical reactions occur that release the alkaline salts, which then wash into freshwater ecosystems.

We called this process freshwater salinization syndrome because it was producing multiple effects on salts, alkalinity and pH, which are fundamental chemical properties of water.



0 400 800 1,200 km



Locations of increasing, decreasing and/or no trends in specific conductance and pH in stream water throughout the continental United States. High-electrical conductivity indicates salinity because salty solutions are full of charged particles that conduct electricity. Credit: Kaushal et al., 2018, CC BY-ND

Different causes by region

Figuring out this process was a team project that required knowledge of limnology (the study of inland waters), geochemistry and geography. The causes vary from one location to another, but the outcomes can be similar.

For example, rivers are becoming more saline and alkaline in parts of North Carolina, Florida, Virginia and other states that use little or no road salt. This is likely due to human-accelerated weathering in locations underlain by limestone (which dissolves when it comes in contact with acid rainwater) and in urbanized areas with lots of concrete infrastructure, as well as urban salt pollution from sewage, water softeners or fertilizers.

Our research was supported by the U.S. National Science Foundation and drew on enormous quantities of monitoring data from ecosystems across the United States collected mainly by the [U.S. Geological Survey](#). We analyzed long-term trends in the chemistry of rivers over five decades and compared these trends across different major river systems and regions.

We also analyzed trends in major estuaries, such as the Hudson River and the Chesapeake Bay, to investigate whether increasingly alkaline inputs from rivers could potentially influence the chemistry of coastal

waters. Our [results](#) show that changes in salts can alter concentrations of pollutants such as excess phosphorus and nutrients that are bound up in sediments at these sites.

Managing salt pollution

Freshwater salinization syndrome is affecting drinking [water supplies](#) in many parts of the United States. In some cases it is [altering the taste of water](#) or [threatening the health](#) of people with hypertension.

There is growing concern that salts in fresh water can [corrode water pipes](#) and release toxic metals such as lead into drinking water. They also can trigger reactions that mobilize other contaminants and pollutants from soils into rivers.

As other scientists have [shown](#), mixtures of salts can be more toxic to aquatic life than just one salt alone. The Environmental Protection Agency does not currently regulate salts as primary contaminants in drinking water, and state and local regulation of salt releases over wide areas from activities such as road treatment are sparse and inconsistent.

We believe there is a serious need for federal regulations and regional plans to [reduce salt pollution in fresh water](#). One strategy would be to reduce use of road salts by calibrating application and adjusting application rates based on temperature. In addition, not all salts are created equal: It may be more efficient to use certain salts as deicers at lower temperatures. Finally, organic de-icing solutions use less [salt](#) than conventional versions.

New forms of water pollution are constantly emerging, and it is important to identify how different human activities accelerate geological processes in nature. Fresh water accounts for only about [3 percent](#) of the Earth's total [water](#) supply (the rest is in the oceans), and

there will always be a need for better understanding and management of this precious resource.

This article was originally published on [The Conversation](#). Read the [original article](#).

Provided by The Conversation

Citation: U.S. rivers are becoming saltier – and it's not just from treating roads in winter (2018, April 17) retrieved 11 May 2024 from <https://phys.org/news/2018-04-rivers-saltier-roads-winter.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.