

Carbon dioxide and iron at the origin of life

May 14 2018

Researchers at the University of Strasbourg have discovered a striking similarity between the way carbon dioxide (CO₂) reacts with metals and the way that ancient microbes use CO₂ to build their biomass, providing potential insight into how chemistry on the early Earth foreshadowed biochemistry in the first organisms.

The study was funded by the European Research Council and is published in the journal *Nature Ecology & Evolution*.

Senior author Dr. Joseph Moran from the Institute for Supramolecular Science and Engineering (ISIS) at the University of Strasbourg and the CNRS said: "Life builds itself from CO₂ and as far as we can tell from biology, it always has. But it's still unclear as to whether that way of doing things extends all the way back to the origin of life, since CO₂ isn't very reactive. One well-explored idea is that more reactive carbon-based molecules such as hydrogen cyanide or formaldehyde were originally used to construct life's molecules. On the other hand, the idea that CO₂ could have been the building block all along is very attractive because it means there might be some continuity between [chemistry](#) on the early Earth and biochemistry as we know it. Much less research has been done in this direction."

To test the hypothesis that CO₂ might have been able to build up biomolecules at the origin of life, the team aimed to mimic the chemistry in one of life's most ancient CO₂ fixation pathways, the acetyl CoA pathway. The enzymes that catalyze the sequence of chemical reactions that make up the acetyl CoA pathway rely heavily on transition

metals such as iron, nickel and cobalt. The team tested how those metals alone react with CO₂ in salty water as a function of temperature and pressure. In the case of iron, the most common of those metals, the main observed organic products were acetate and pyruvate—the same intermediates and end-products of the acetyl CoA pathway.

Dr. Moran said: "The similarity between the chemistry happening on the surface of the metal and the chemistry happening in this [biological pathway](#) is remarkable. It suggests there may indeed be some continuity between prebiotic chemistry and the biochemistry of early life. In these cases, enzymes may have simply served to speed up and control existing chemical reactions."

One important difference between the biological pathway and the discovered chemistry is that the former uses hydrogen gas as its source of electrons, whereas the latter ultimately takes its electrons from the [metal](#). "This might be a clue that very early life got its electrons elsewhere before the complex electron bifurcation mechanisms used by modern [life](#) had evolved to exploit hydrogen gas."

The paper, "Native iron reduces CO₂ to intermediates and end-products of the acetyl CoA [pathway](#)," is published in the journal *Nature Ecology & Evolution*.

More information: Sreejith J. Varma et al. Native iron reduces CO₂ to intermediates and end-products of the acetyl-CoA pathway, *Nature Ecology & Evolution* (2018). DOI: 10.1038/s41559-018-0542-2 , doi.org/10.1038/s41559-018-0542-2

Provided by University of Strasbourg

Citation: Carbon dioxide and iron at the origin of life (2018, May 14) retrieved 27 April 2024 from <https://phys.org/news/2018-05-carbon-dioxide-iron-life.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.