

# Copepod mortality rates have been linked to jellyfish abundance and warmer, windier conditions

May 1 2018

---



*Calanus helgolandicus*. Credit: Plymouth Marine Laboratory

Zooplankton may be tiny, but their importance to the health of the planet is not – they are microscopic life that makes a huge contribution to the

fine balance of ocean ecosystems. Copepods make up a significant proportion of zooplankton biomass – tiny invertebrates that are the key prey resource for many marine animals, from other zooplankton and fish larvae to sharks and whales, and are a crucial part of the food web.

Mortality—be it death as a result of environmental conditions or by becoming a meal for a larger predator—is of course an integral part of zooplankton [population](#) dynamics. But these losses are generally harder to quantify than the gains to a population via reproduction. PML scientists studying the important copepod *Calanus helgolandicus* have now discovered links between their [mortality](#) rates and predators, and also temperature and wind speed, and how these factors influence the delicate balance of [copepod](#) populations.

Warmer waters could mean an increase in danger for adult copepods, in the form of more abundant gelatinous predators like jellyfish and comb jellies. PML researchers have found that higher mortality rates are related to both higher temperatures and greater predator abundance, with gelatinous predators highlighted as the leading agent for *C. helgolandicus* population control in the English Channel. Windy weather, too, has an impact. For copepods that manage to evade predators, environmental conditions can still take their toll; deaths that don't involve becoming a [predator](#)'s lunch are instead correlated with high wind speeds and greater ocean turbulence.

"We currently know very little about zooplankton death, basically because we can't tag them to find out where they go and how they die," said lead author Jacqueline Maud. "Our work goes some way towards identifying and quantifying the different types of mortality, so that we can understand what's really going on in marine ecosystems. In terms of *C. helgolandicus*, we found that it is the jellies that really control the population, and also that extreme weather events can impact on this tiny—but incredibly important—member of the [zooplankton](#)

community."

**More information:** Jacqueline L. Maud et al, Mortality of *Calanus helgolandicus* : Sources, differences between the sexes and consumptive and nonconsumptive processes, *Limnology and Oceanography* (2018).  
[DOI: 10.1002/lno.10805](https://doi.org/10.1002/lno.10805)

Provided by Plymouth Marine Laboratory

Citation: Copepod mortality rates have been linked to jellyfish abundance and warmer, windier conditions (2018, May 1) retrieved 14 August 2024 from <https://phys.org/news/2018-05-copepod-mortality-linked-jellyfish-abundance.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.