

If El Ninos happen twice as often in the future, what happens to seabirds?

May 7 2018



Brandt's cormorant. Credit: Annie Schmidt/UC Davis

More frequent El Niño events in the future may have surprising impacts on seabirds and some fish species, according to a study from the University of California, Davis.

El Niños are unusually warm ocean conditions that occur every two to seven years off the Pacific Coast, bringing with them poor ocean productivity and sometimes catastrophic weather conditions. Fossil coral records and [climate change](#) models indicate that El Niños occurred both more and less frequently over the past 1,000 years than they do now, and climate change may speed up or slow down their frequency in the future.

In a modeling study recently published in the journal *Theoretical Ecology*, UC Davis researchers in the Department of Wildlife, Fish and Conservation Biology wondered how changes in frequency of El Niño and its more favorable, cool-water counterpart La Niña might affect Brandt's cormorant. The seabird was selected as a model species because of its known sensitivity to environmental changes.

"We expected that if you increased the frequency of El Niños it would have a negative impact on the population," said lead author Annie Schmidt, a Ph.D. student at UC Davis during the time of the study and currently a researcher at the nonprofit Point Blue Conservation Science. "It turns out it was exactly the opposite."

Taking the good with the bad

The study's models indicated that doubling the frequency of the El Niño-Southern Oscillation (ENSO) phenomenon, which includes El Niño and La Niña, unexpectedly resulted in higher population numbers and a lower chance of extinction for Brandt's cormorants.



Pair of Brandt's cormorants on the Farallon Islands. Credit: Annie Schmidt/UC

Davis

That's because ENSO alternates between unfavorably warm El Niño conditions followed by favorable, cooler La Niña phases. Shorter, more frequent El Niños may mean the birds experience shorter periods of poor conditions, with La Niña following to help them bounce back.

"If we speed things up, then we'll get shorter bad and shorter good periods," said co-author Louis Botsford, a professor in the UC Davis Department of Wildlife, Fish and Conservation Biology. "If we slow things down, we get longer good and longer bad periods. The latter is not good for this population because populations are driven to low levels and go extinct when you hold them down for long periods of time."

By interspersing the failures with the successes, the birds are better able to cope if El Niños become more frequent.

The study applies to other species, as well. Brandt's cormorants can live for more than 20 years. Species with similar life histories would likely respond the same way, the authors said. This could include other long-lived seabirds and some fish, such as Pacific hake and some rockfish.

Climate change's effects on seabirds

Does this study indicate that climate change could be beneficial for Brandt's cormorants?



Colony of Brandt's cormorants on the Farallon Islands, off the coast of San Francisco. Credit: Annie Schmidt/UC Davis

"I wouldn't go that far," Schmidt said. "This study is looking at a narrow scope of potential change. In terms of overall effects of climate change, there are a lot of other factors that come into play with this species and whole ecosystems that can be cumulative."

What the study does indicate is the importance of understanding how different species respond to different frequencies of change in the environment. That is seldom considered when calculating the probability of extinction.

Future uncertain

The authors also point out that their study is not predictive. Exactly how El Niño frequency may be altered by climate change is uncertain. If the nature of the oscillation were to change, with unfavorable El Niño conditions becoming more frequent or longer, while favorable La Nina conditions did not change, the results could be very different.

Additional authors include John Eadie and Patrick Kilduff of UC Davis, and Russell Bradley and Jaime Jahncke of Point Blue Conservation Science.

More information: Annie E. Schmidt et al, Changing environmental spectra influence age-structured populations: increasing ENSO frequency could diminish variance and extinction risk in long-lived seabirds, *Theoretical Ecology* (2018). [DOI: 10.1007/s12080-018-0372-5](https://doi.org/10.1007/s12080-018-0372-5)

Provided by UC Davis

Citation: If El Ninos happen twice as often in the future, what happens to seabirds? (2018, May 7) retrieved 20 March 2024 from <https://phys.org/news/2018-05-el-ninos-future-seabirds.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--