

## New model explains what we see when a massive black hole devours a star

May 29 2018, by Tim Stephens



n the figure we see a cross section of what happens when the material from the disrupted star is devoured by the black hole. An accretion disk is formed (disk) by the material. There is too much material for it to pass into the black hole at



once. It is heated up in the process and emits vast amounts of light and radiation, visible from Earth (Double arrow). Dr. Jane Dai's computer model takes the difference in viewing angle from Earth into account, which means we are now able to categorize the variations in observations correctly. This means we can study the properties of the black hole, and learn about a celestial body we would otherwise not be able to see. Credit: Niels Bohr Institute

A star that wanders too close to the supermassive black hole in the center of its galaxy will be torn apart by the black hole's gravity in a violent cataclysm called a tidal disruption event (TDE), producing a bright flare of radiation. A new study led by theoretical astrophysicists at the University of Copenhagen's Niels Bohr Institute and UC Santa Cruz provides a unified model that explains recent observations of these extreme events.

The breakthrough study, published in *Astrophysical Journal Letters*, provides a new theoretical perspective for a fast-growing research field.

"Only in the last decade or so have we been able to distinguish TDEs from other galactic phenomena, and the new model will provide us with the basic framework for understanding these rare events," said coauthor Enrico Ramirez-Ruiz, professor and chair of astronomy and astrophysics at UC Santa Cruz and Niels Bohr Professor at the University of Copenhagen.

In most galaxies, the central black hole is quiescent, not actively consuming any material and therefore not emitting any light. Tidal disruption events are rare, only happening about once every 10,000 years in a typical galaxy. When an unlucky star gets torn apart, however, the black hole is "overfed" with stellar debris for a while and emits intense radiation.



"It is interesting to see how materials get their way into the black hole under such extreme conditions," said first author Jane Lixin Dai, assistant professor at the University of Copenhagen, who led the study. "As the black hole is eating the stellar gas, a vast amount of radiation is emitted. The radiation is what we can observe, and using it we can understand the physics and calculate the black hole properties. This makes it extremely interesting to go hunting for tidal disruption events."

While the same physics is expected to happen in all tidal disruption events, about two dozen of which have been observed so far, the observed properties of these events have shown great variation. Some emit mostly x-rays, while others emit mostly visible and ultraviolet light. Theorists have been struggling to understand this diversity and assemble different pieces of the puzzle into a coherent model.

## Viewing angle

In the new model, it is the viewing angle of the observer that accounts for differences in the observations. Galaxies are oriented randomly with respect to the line of sight of observers on Earth, who see different aspects of a tidal disruption event depending on its orientation.

"It is like there is a veil that covers part of a beast," Ramirez-Ruiz explained. "From some angles we see an exposed beast, but from other angles we see a covered beast. The beast is the same, but our perceptions are different."

The model developed by Dai and her collaborators combines elements from general relativity, magnetic fields, radiation, and gas hydrodynamics. It shows what astronomers can expect to see when viewing tidal disruption events from different angles, allowing researchers to fit different events into a coherent framework.



Survey projects planned for the next few years are expected to provide much more data on tidal disruption events and will help greatly expand this field of research, according to Dai. These include the Young Supernova Experiment (YSE) transient survey, led by the DARK Cosmology Centre at the Niels Bohr Institute and UC Santa Cruz, and the Large Synoptic Survey Telescopes being built in Chile.

"We will observe hundreds to thousands of tidal <u>disruption</u> events in a few years. This will give us a lot of 'laboratories' to test our model and use it to understand more about <u>black holes</u>," Dai said.

In addition to Dai and Ramirez-Ruiz, the coauthors include Jonathan McKinney, Nathaniel Roth, and Cole Miller at the University of Maryland, College Park. State-of-the-art computational tools were employed to solve the puzzle, and the simulations were carried out by Dai and Roth on the recently acquired large computer cluster made possible by a grant from the Villum Foundation for Jens Hjorth, head of DARK Cosmology Centre, as well as clusters funded by the U.S. National Science Foundation and NASA.

**More information:** Lixin Dai et al. A Unified Model for Tidal Disruption Events, *The Astrophysical Journal* (2018). DOI: 10.3847/2041-8213/aab429, iopscience.iop.org/article/10. ... 041-8213/aab429/meta

## Provided by University of California - Santa Cruz

Citation: New model explains what we see when a massive black hole devours a star (2018, May 29) retrieved 3 May 2024 from <u>https://phys.org/news/2018-05-massive-black-hole-devours-star.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.