

Morphing twisted nanoscale objects to tailor applications in future technologies

May 3 2018, by Chris Melvin



An impression of a chiral molecule moving through various configurations as it transitions from one handedness to another. Credit: Ventsislav Valev and Joel Collins



For the first time scientists have created a way to model the interaction between light and twisted molecules, as these molecules transition from left- to right-handed versions, or vice versa. The transitional forms offer a deeper insight into material symmetries and their unexpected behaviour could lead to improved design of telecoms components.

Many <u>molecules</u>, including important pharmaceuticals and valuable chemicals, exist in two "chiral" forms - they have the same chemical structure arranged in mirror images, termed left-handed and righthanded forms. This can alter their properties and is therefore important to fully understand how the compound interacts with other molecules, or light.

Typically, it has only been possible to study either the left- or righthanded chiral form but nothing in between, however ideally scientists would like to gradually morph a shape from one handedness to the other and observe how the effects of this change translate into physical properties.

Now a research team from the Department of Physics at the University of Bath, working with colleagues at University College London, Belgium and China, has created a way to do exactly that.

Their unique method involves manufacturing metallic nano-scale "artificial molecules" representative of 35 intermediate stages along the way of a geometric transformation, from one handedness to the other. At this nano-scale, the shape of the artificial molecule affects its optical properties, so by using twisted laser light the team studied the properties of the various stages, as the artificial molecules morphed from left to right handedness.

PhD student Joel Collins said: "We were able to follow the properties of a chiral artificial molecule, as it was morphed from left- to right-handed



form, through two different routes. No-one has done this before. Surprisingly, we found that each route leads to a different behavior.

"We measured the difference in absorption of left and right circularly polarized light, known as circular-dichroism (CD). Along one route, the artificial molecules behaves as might be expected, with progressively decreasing CD, and eventually a reversal of the CD, for the mirrored structure. However, along the second route, the CD reversed several times, even before the structure changed handedness."

The research is published in the journal Advanced Optical Materials.

Dr Ventsislav Valev who led the research said: "This is actually a very elegant idea but it has only become a possibility thanks to the recent advances in nanofabrication.

"In chemistry, you can't tune the twist of a <u>chiral molecule</u>, so every scientist who studies such molecules needs to tune the wavelength of <u>light</u>. We have demonstrated a new, complementary physical effect, where we fix the wavelength and tune the twist of the chiral artificial molecule. In many cases, our approach is more practical; for instance, when we're designing telecoms components, where the optical wavelength is pre-determined."

More information: Joel T. Collins et al, Enantiomorphing Chiral Plasmonic Nanostructures: A Counterintuitive Sign Reversal of the Nonlinear Circular Dichroism, *Advanced Optical Materials* (2018). DOI: 10.1002/adom.201800153

Provided by University of Bath



Citation: Morphing twisted nanoscale objects to tailor applications in future technologies (2018, May 3) retrieved 2 May 2024 from https://phys.org/news/2018-05-morphing-nanoscale-tailor-applications-future.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.