


Polarized light—a simple route to highly chiral materials

May 9 2018



Nanostructures prepared by RCP and LCP light irradiation. Credit: 2018 Tetsu Tatsuma, Institute of Industrial Science, The University of Tokyo

Researchers at the University of Tokyo used an efficient method to create chiral materials using circularly polarized light. Depending on whether left- or right-polarized, the light source induced electric fields at opposite corners of gold nanocuboids on TiO_2 . Via plasmon-induced charge separation, the gold converted Pb^{2+} into PbO_2 tips deposited at the corners, resulting in a chiral plasmonic nanostructure with high enantiomeric excess. Materials with such a chiral form are useful for sensing and asymmetric synthesis.

Chirality is at the heart of chemical research and much technology. For organic chemists, choosing between the left- and right-handed isomers

of molecules is all part of a day's work. However, many solid [materials](#) also have enantiomeric forms, giving rise to a range of applications.

Organic chemists generally rely on an arsenal of laboratory reactions to control chiral purity. For materials, there is another, more elegant approach—circularly polarized [light](#), which is readily made, and can be either left-circularly polarized (LCP) or right-circularly polarized (RCP). In material synthesis, the opposite twists of LCP and RCP light indirectly lead to structures that are mirror images of each other.

Previously, this strategy has been hampered in practice. Now, researchers at The University of Tokyo's Institute of Industrial Science have successfully created chiral nanostructures from particles of gold (Au). The trick was to use circularly polarized light to generate electric fields, which localize differently depending on LCP or RCP. This in turn drove the chiral deposition of a dielectric material.

As described in a study reported in *Nano Letters*, the researchers first deposited Au nanocuboids—essentially miniature rectangular gold bars—on a TiO₂ substrate.

As study co-author Koichiro Saito explains, "Under a beam of [circularly polarized light](#), electric fields built up around the cuboids—but at one pair of corners for LCP rotation, and the opposite pair under RCP light. At this point, we had achieved chirality, but in electric rather than material form."

The chirality of the electric field was then transferred to the material itself by plasmon-induced charge separation, in which Pb²⁺ ions were oxidized through the chirally distributed electric fields. This deposited PbO₂, a dielectric material, at either one set of cuboid corners or the other, depending on the original light source. Electron microscopy showed the gold bars transformed into non-superimposable mirror

images, the hallmark of chirality.

"This is the first time a chiral material has been made by exploiting plasmon resonance," co-author Tetsu Tatsuma says. "No other source of chirality is needed but light itself. Nanoscale chiral plasmonic materials are highly useful for sensing and asymmetric synthesis, and our process makes them much more efficient to produce. Plus, we don't think it's limited to one product—other chiral nanomaterials have an incredible range of functions in modern technology."

More information: Koichiro Saito et al, Chiral Plasmonic Nanostructures Fabricated by Circularly Polarized Light, *Nano Letters* (2018). [DOI: 10.1021/acs.nanolett.8b00929](https://doi.org/10.1021/acs.nanolett.8b00929)

Provided by University of Tokyo

Citation: Polarized light—a simple route to highly chiral materials (2018, May 9) retrieved 24 April 2024 from <https://phys.org/news/2018-05-polarized-light-a-simple-route-highly.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.