

Study shows sea turtle nesting beaches threatened by microplastic pollution

May 1 2018, by Zachary Boehm



Student researcher Victoria Beckwith surveyed 10 important loggerhead turtle nesting beaches along the Gulf coast. Microplastics were present at each site. Credit: Victoria Beckwith

Plastic is famous for its unyielding durability, making it perfect for consumer products but a unique and persistent menace to the natural environment.

For the loggerhead sea turtles that nest on the once-pristine beaches bounding the Gulf of Mexico, millimeters-thick pieces of broken down plastic—called microplastics—pose a particularly urgent threat.

A new study from Florida State University researchers shows that increasing [microplastic](#) accumulation along the Gulf's beaches could alter the composition of shoreline sand and jeopardize the turtles' sensitive incubation environments.

Their findings were published in the journal *Marine Pollution Bulletin*.

"With increasing populations, higher demand for resources and more use of plastic, we're having a lot more plastic and microplastic appearing as marine debris," said the study's coauthor Mariana Fuentes, assistant professor of Earth, Ocean and Atmospheric Science (EOAS). "In these coastal areas, we're seeing significantly more pollution."

For the study, EOAS student researcher Valencia Beckwith surveyed the Northern Gulf of Mexico Loggerhead Recovery Unit's 10 most important loggerhead turtle nesting sites in Florida.

Sand samples collected throughout the region revealed that microplastics were present at every site. More alarming, the highest concentrations of microplastics were found consistently in the dunes, where sea turtles tend to nest.



Loggerhead hatchlings. Credit: Wikimedia Commons

Plastic has a tendency to retain large amounts of heat in response to comparably moderate increases in [temperature](#). If enough plastic is present in a sandy environment, the area could experience measurable temperature increases.

This dynamic is of particular concern in sea turtle nests, Fuentes said. For marine turtle eggs, incubation temperature is destiny.

"Sea turtles have temperature dependent sex determination, which means their sex is determined by the sand temperature," Fuentes said. "Changes in incubation temperatures might modify the sex ratios produced on these nesting beaches, but at this stage we don't know how much microplastic is needed to see those changes."

In subsequent research, Beckwith and Fuentes plan to expand upon these findings and investigate the specific ways that microplastic might alter the temperature profile of the sediment on important nesting beaches.

"The first step was to see whether sea [turtles](#) are exposed to microplastics," she said. "Next we'll explore its potential impacts."

Earth's oceans have long been blighted by pollution, and vulnerable species like [sea turtles](#) have borne the brunt of decades of irresponsible waste. But Fuentes remains optimistic about the future. She said that shifting attitudes could translate into positive changes in policy and behavior.

"There is a lot of hope," Fuentes said. "We're beginning to see more and more initiatives providing incentives to discourage the use of plastics. I see my students making those changes every day. It's up to everyone."

More information: Valencia K. Beckwith et al, Microplastic at nesting grounds used by the northern Gulf of Mexico loggerhead recovery unit, *Marine Pollution Bulletin* (2018). [DOI: 10.1016/j.marpolbul.2018.04.001](https://doi.org/10.1016/j.marpolbul.2018.04.001)

Provided by Florida State University

Citation: Study shows sea turtle nesting beaches threatened by microplastic pollution (2018, May 1) retrieved 6 May 2024 from <https://phys.org/news/2018-05-sea-turtle-beaches-threatened-microplastic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.