

Disease afflicting frogs becoming deadlier

June 26 2018



Pumpkin Toadlet, endemic to Brazil's Atlantic forest, is highly susceptible to the hybrid Bd fungus, according to UA-led research. Credit: Dr. Gui Becker

A disease-inducing fungus in amphibians worldwide could become deadlier as different genetic variations emerge, according to research led by The University of Alabama.

Biologists tested the harmfulness of a Brazilian hybrid form of *Batrachochytrium dendrobatidis*, known simply as Bd or the [amphibian chytrid fungus](#), as well as both parent forms, on species of frogs from the Brazilian Atlantic Forest. They found infections and strength of the illness increased with the hybrid form.

Published today in *Scientific Reports*, the research is the first to demonstrate hybrids of the [fungus](#) can be deadlier than more common and widespread fungal genotypes.

"With globalization facilitating pathogen spread across continents, hybridization of chytrids might lead to new epidemic waves reducing [amphibian](#) biodiversity in both tropical and temperate regions," said Dr. Gui Becker, UA assistant professor of biological sciences.

Becker was a co-author on the paper, "Hybrids of amphibian chytrid show high virulence in native hosts," which was led by Dr. Sasha Greenspan, a UA post-doctoral researcher.

When the paper was written a few months ago, the Brazilian hybrid of the fungus was the only one documented, but new evidence shows another hybrid of the fungus emerged in South Africa, Greenspan said.

"The genetics of this fungus are changing, and with these changes may come new risks for frog populations," she said.

Infectious outbreaks of the disease caused by the fungus, chytridiomycosis, have afflicted amphibians for at least 40 years on every continent except Antarctica, sometimes killing massive numbers

of animals and, occasionally, extinguishing some species.

The fungus' complex genetic history is not completely understood, but a relatively new version, on an evolutionary timescale, is known as the Global Pandemic Lineage, or Bd-GPL. What led to its emergence is unknown, but it is particularly deadly.

In Brazil, Bd-GPL reproduced with another lineage, a process known as hybridization, and the UA-led research shows the hybrid fungus can be even more deadly than Bd-GPL in native frogs, Greenspan said.

"This is significant because it shows that hybrids of this fungus can increase disease risk," she said.

The team conducted the research at the University of Campinas in the state of São Paulo, Brazil, and included a diverse array of frogs native to the Brazilian Atlantic Forest, including *Brachycephalus ephippium*, called a pumpkin toadlet among other names; *Ischnocnema parva*; and *Dendropsophus minutus*, a lesser tree [frog](#).

"Since it is well-established that Bd has a dynamic genome and is now nearly globally distributed, this finding highlights the importance of global surveillance to detect future genomic shifts in this disease system that could lead to new outbreaks of chytridiomycosis," Greenspan writes in the paper.

More information: S. E. Greenspan et al. Hybrids of amphibian chytrid show high virulence in native hosts, *Scientific Reports* (2018). [DOI: 10.1038/s41598-018-27828-w](https://doi.org/10.1038/s41598-018-27828-w)

Provided by University of Alabama in Tuscaloosa

Citation: Disease afflicting frogs becoming deadlier (2018, June 26) retrieved 2 May 2024 from <https://phys.org/news/2018-06-disease-afflicting-frogs-deadlier.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.