


# Outflowing gas from galaxy supermassive black hole nuclei

June 4 2018

---



An image of the galaxy Markarian 348 in the ultraviolet. Its active nucleus powers outflowing atomic gas, and new observations of it and four similar galaxies have been able to image the outflow as well as a rotating gas component.  
Credit: NASA/GALEX

Supermassive black holes at the nuclei of most galaxies, including our

Milky Way, develop gradually as material accretes onto the seed black hole. The physical processes that drive this growth – the so-called feeding and feedback processes – occur in the vicinity of the galaxy nucleus. When the accretion becomes active, radiation is emitted that illuminates and ionizes the gas in the vicinity of the nucleus.

Accretion disc winds can interact with the gas to produce outflowing gas that is observed to reach velocities of hundreds of km/sec. Relativistic jets of particles emanating from the black hole can also interact with his material. These various kinds of feedback are essential to avoid producing overly massive galaxies.

Clear evidence for all these processes has been detected in their optical emission lines of ionized atoms, whose velocities can be measured. However it has been much hard to obtain spatial information about the geometry of the excited gas. CfA astronomer Martin Elvis and nine colleagues used the Gemini eight-meter telescope and a powerful new instrument that records both high-resolution spatial (as small as a few hundred light-years in size) and [velocity](#) information.

The team studied five relatively nearby galaxies known to have active black hole nuclei with bright atomic emission. They discovered that in all cases the gas has two major components, one rotating and one outflowing. But otherwise the [galaxies](#) are all somewhat different: in one the gas rotates opposite to its stars, in another only one lobe of the outflow can be seen, and there are other differences as well. The new paper is just the first in a series expected to probe and model in detail how nuclear [black holes](#) grow.

**More information:** I C Freitas et al. Outflows in the narrow-line region of bright Seyfert galaxies – I. GMOS-IFU data, *Monthly Notices of the Royal Astronomical Society* (2018). [DOI: 10.1093/mnras/sty303](https://doi.org/10.1093/mnras/sty303)

Provided by Harvard-Smithsonian Center for Astrophysics

Citation: Outflowing gas from galaxy supermassive black hole nuclei (2018, June 4) retrieved 3 May 2024 from <https://phys.org/news/2018-06-outflowing-gas-galaxy-supermassive-black.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.