

Study: Populations of widely spread tree species respond differently to climate change

June 8 2018



The study examined growth rings from the world's southernmost conifer tree species, *Pilgerodendron uviferum*. Credit: Andrés Holz, Portland State University

A new Portland State University study shows that not all populations of a single, widely spread tree species respond the same to climate change, something scientists will need to consider when making climate change projections.

The study, published online recently in the *Journal of Biogeography*, examined [growth](#) rings from the world's southernmost conifer tree species, *Pilgerodendron uviferum*, to see if its growth rate responded

differently to [climate](#) change over time and depending on its location in northern, central or southern Patagonia.

The study, led by PSU geography professor Andrés Holz, found that during the first half of the 20th century, the climate-growth relationships were relatively similar between the southern (poleward) and northern (equatorward) range edges, but diverged after the 1950s when warming in southern South America increased.

In the northern and southern edges, where it can be relatively dry during the spring and summer months, wetter-than-average conditions resulted in increased tree growth. However, in central Patagonia, the region that gets the most rain, growth was mostly dependent on drier and sunnier conditions.

But in the 1950s, warmer conditions globally as well as an upward trend in the [climate phenomena](#) known as the Antarctic Oscillation, or Southern Annular Mode, shifted storm tracks poleward resulting in a decline in precipitation in northern Patagonia and an increase in precipitation in central and southern Patagonia. As a result, trees in southern Patagonia reversed trend and now their growth is favored by drier, sunnier conditions.

"We found that not only can we see different populations of a widely distributed species respond differently to natural [climate variability](#) in a given region, but also over time within each of these regions and because climate variability is amplified by climate change, their tree growth responses to climate can change from a bit to a lot," said Holz, who also serves as a faculty fellow in PSU's Institute for Sustainable Solutions.

Holz said the findings suggest that when trying to simulate how different species and ecosystems are going to respond to climate change in the next 10, 50 or 100 years, a single parameter per species, and even per

population, cannot be used to represent an entire [species](#)' growth.

"We show that that would not work because they respond differently over time and space," Holz said.

More information: Andrés Holz et al, Radial growth response to climate change along the latitudinal range of the world's southernmost conifer in southern South America, *Journal of Biogeography* (2018).

[DOI: 10.1111/jbi.13199](https://doi.org/10.1111/jbi.13199)

Provided by Portland State University

Citation: Study: Populations of widely spread tree species respond differently to climate change (2018, June 8) retrieved 24 April 2024 from <https://phys.org/news/2018-06-populations-widely-tree-species-differently.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--