

When the river runs high

June 15 2018



River Kvirila at Sachkhere, Georgia. Credit: Wikipedia

A massive world-wide study of dry riverbeds has found they're contributing more carbon emissions than previously thought, and this could help scientists better understand how to fight climate change.



Dr. Nathan Waltham from the Tropical Water and Aquatic Eco-systems Research Centre (TropWATER) and James Cook University in Australia, joined scientists from 22 other countries who looked at 212 dry riverbeds on every continent on earth.

He said the contribution of intermittent rivers and <u>streams</u> to the process of carbon cycling—the process by which carbon is circulated around the ecosystem—is largely ignored.

"There is a substantial amount of plant litter that accumulates in dry riverbeds and when they flow again this material can breakdown rapidly. We've now estimated the potential short-term CO2 emissions during these rewetting events."

"We believe that a single pulse of CO2 <u>emission</u> upon litter rewetting contributes up to 10% of the daily CO2 emission compared to perennial rivers and streams, particularly in temperate climates. What this means is that the contributions of intermittent rivers and streams should be included in global carbon-cycling assessments," said Dr. Waltham.

The scientists found that aridity, surrounding vegetation, channel width and dry-phase duration explained most variability in the quantity and decomposability of <u>plant litter</u>.

He said that the new data shows the contribution of CO2 from intermittent rivers and streams is higher than previously thought.

"Taking rivers and streams that only flow at certain times into account would improve estimates of the consequences of global <u>climate change</u> on carbon cycling—given that the extent of these rivers and streams will increase, and periods of drying will become more prolonged in many regions," said Dr. Waltham.



About the research

Intermittent rivers, as the name suggests, sometimes stop flowing and can dry completely. Although far less studied than permanent rivers, they could represent half of the world's river network and, in response to climate change and increasing water demands, may come to dominate the landscape in some regions.

The results of this global study have been published in *Nature Geosciences*.

More information: T. Datry et al, A global analysis of terrestrial plant litter dynamics in non-perennial waterways, *Nature Geoscience* (2018). DOI: 10.1038/s41561-018-0134-4

Provided by James Cook University

Citation: When the river runs high (2018, June 15) retrieved 18 April 2024 from https://phys.org/news/2018-06-river-high.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.