



Supramolecular complex formation—anthracene macrocycle and C60 fullerene

June 8 2018



Nano-Saturn



Credit: Wiley

Saturn is the second largest planet in our solar system and has a characteristic ring. Japanese researchers have now synthesized a molecular "nano-Saturn." As the scientists report in the journal

Angewandte Chemie, it consists of a spherical C₆₀ fullerene as the planet and a flat macrocycle made of six anthracene units as the ring. The structure is confirmed by spectroscopic and X-ray analyses.

Nano-Saturn systems with a spherical molecule and a macrocyclic ring have been a fascinating structural motif for researchers. The ring must have a rigid, circular form, and must hold the molecular sphere firmly in its midst. Fullerenes are ideal candidates for the nano-sphere. They are made of carbon atoms linked into a network of rings that form a hollow sphere. The most famous [fullerene](#), C₆₀, consists of 60 [carbon atoms](#) arranged into 5- and 6-membered rings like the leather patches of a classic soccer ball. The electrons in their double bonds, known as the π -electrons, are in a kind of "[electron cloud](#)," able to freely move about and have binding interactions with other molecules, such as a macrocycle that also has a "cloud" of π -electrons. The attractive interactions between the electron [clouds](#) allow fullerenes to lodge in the cavities of such macrocycles.

A series of such complexes has previously been synthesized. Because of the positions of the electron clouds around the macrocycles, it was previously only possible to make rings that surround the fullerene like a belt or a tire. The ring around Saturn, however, is not like a "belt" or "tire," it is a very flat disc. Researchers working at the Tokyo Institute of Technology and Okayama University of Science (Japan) wanted to properly imitate this at nanoscale.

Their success resulted from a different type of bonding between the "nano-planet" and its "nano-ring." Instead of using the attraction between the π -electron clouds of the fullerene and macrocycle, the team working with Shinji Toyota used the weak attractive interactions between the π -electron cloud of the fullerene and non- π -electron of the carbon-hydrogen groups of the macrocycle.

To construct their "Saturn [ring](#)," the researchers chose to use anthracene units, molecules made of three aromatic six-membered carbon rings linked along their edges. They linked six of these units into a macrocycle whose cavity was the perfect size and shape for a C₆₀ fullerene. Eighteen hydrogen atoms of the macrocycle project into the middle of the cavity. In total, their interactions with the fullerene are enough to give the complex enough stability, as shown by computer simulations. By using X-ray analysis and NMR spectroscopy, the team was able to prove experimentally that they had produced Saturn-shaped complexes.

More information: Yuta Yamamoto et al. Nano-Saturn: Experimental Evidence of Complex Formation of an Anthracene Cyclic Ring with C₆₀, *Angewandte Chemie International Edition* (2018). [DOI: 10.1002/anie.201804430](https://doi.org/10.1002/anie.201804430)

Provided by Wiley

Citation: Supramolecular complex formation—anthracene macrocycle and C₆₀ fullerene (2018, June 8) retrieved 7 May 2024 from <https://phys.org/news/2018-06-supramolecular-complex-formationanthracene-macrocycle-c60.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--