

Toxic leftovers from gold mine found in snowshoe hares

June 14 2018, by Som Niyogi And Solomon Amuno



Snowshoe hares near the now closed Giant Mine outside of Yellowknife, N.W.T show signs of arsenic contamination. Credit: <u>Denali NPS/flickr</u>, <u>CC BY-SA</u>

Even though it was closed decades ago, the Giant Mine on the outskirts of Yellowknife has left a long environmental legacy.



The gold extraction process, which required roasting ores at extremely high temperatures, created a toxic byproduct called <u>arsenic trioxide</u>. For about 55 years (1948-2004), <u>arsenic</u> and other toxic elements were released into the environment, causing <u>widespread contamination of the terrestrial and aquatic ecosystems around Yellowknife</u>.

About <u>237,000 tonnes of arsenic trioxide dust is buried</u> underground, and <u>several nearby lakes show arsenic contamination</u>.

Elevated <u>arsenic levels</u> have also been reported in soil, vegetation and fish around Yellowknife, but we knew little about how it has affected the health of the <u>small mammals</u> that live in the area.

Many of these fur-bearing animals are still being trapped for their pelts and for food, so knowing their arsenic levels is also important for human health.

Weak bones

Small mammals can serve as sentinels for environmental contamination. Snowshoe hares (*Lepus americanus*) live in a relatively small area and eat soil, so they are likely to accumulate higher levels of arsenic and other trace metals from the environment.

Exposure to elevated levels of arsenic can cause damage to the liver and other organs. And cadmium, a toxic metal and another byproduct of the gold extraction process, can replace calcium in the bones, leading to bone deformities and weakness.

In humans, chronic arsenic exposure (usually from water) can lead to changes in skin colour, skin growths and cancers of the skin, lung and internal organs.



When we measured arsenic and <u>cadmium levels</u> in hares living within two kilometres of the Giant Mine and compared them to hares living about 20 kilometres away from Yellowknife, the results were striking.

The arsenic levels in the guts of snowshoe hares living near the Giant Mine were 20-50 times greater than those living away from it. We also saw higher concentrations of arsenic in the organs and nails of the Giant Mine hares.

Cadmium levels were also higher but the difference wasn't as marked. Hares from both locations had weaker bones and showed signs of osteoporosis, probably due to <u>chronic exposure to cadmium</u>.

Ecological implications

This chronic exposure to elevated levels of arsenic and cadmium may explain why snowshoe hares living near the Giant Mine are in poor health.

Wildlife living in metal contaminated areas in other parts of the world have also shown problems with reproduction, osteoporosis, neurological damage and chronic metabolic disease. But in Canada, it's the first time we've seen small wild mammals with chronic arsenic poisoning.

The high levels of pollutants could compromise the long-term survival of the snowshoe <u>hare</u> and other small mammals in the Yellowknife area.

The high arsenic and <u>cadmium</u> burden in hares could have consequences for other animals that prey on them, such as foxes, wolves or other carnivorous mammals, and <u>for the people who hunt them</u>.

This article was originally published on <u>The Conversation</u>. Read the <u>original article</u>.



Provided by The Conversation

Citation: Toxic leftovers from gold mine found in snowshoe hares (2018, June 14) retrieved 17 July 2024 from <u>https://phys.org/news/2018-06-toxic-leftovers-gold-snowshoe-hares.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.