

Diverse salmon populations enable 'resource surfing' bears to eat tons of fish

July 23 2018



Coho salmon, Tillamook State Forest, Oregon. Credit: Oregon Department of Forestry

Research shows that Kodiak brown bears that sync their stream-to-stream movements to salmon spawning patterns eat longer and more than

bears that don't, with one bear in the study consuming greater than 2 tons of fish in one summer.

Individual sockeye salmon populations spawn for about 40 days, but "resource surfing" bears can fish for three times that long, biologists have learned.

Worldwide, prolonged salmon availability is increasingly under threat from hatchery supplementation that tends to reduce the genetic diversity underpinning different spawning times. In addition, bears' ability to follow salmon waves is hampered by industrial development such as mining.

Findings were just published in *Scientific Reports*.

"This study is the first to link actual metrics of bear consumption to their foraging behavior and movements," said co-author Jonathan Armstrong, assistant professor of fisheries and wildlife in the Oregon State University College of Agricultural Sciences.

Armstrong collaborated with corresponding author Will Deacy and scientists from the University of Montana, Washington State University and the Kodiak National Wildlife Refuge to collar 33 female bears and track them for a year over a 1-million-square-kilometer portion of Kodiak Island.

At the end of those 12 months, the team recaptured 18 of the bears and took hair samples that they measured for mercury. Salmon absorb mercury from what they eat in the ocean, and the amount of mercury in a bear's hair indicates how much fish it dined on—the more mercury, the more salmon it ate.

"Salmon consumption ranged from around 300 kilograms for one bear

up to almost 2,000 kilograms for the biggest salmon eater," said Deacy, a postdoctoral scholar at OSU. "This study complements our other research to show how bears depend on diverse salmon populations."

On average, the bears in the study ate more than 1,000 kilograms of salmon apiece, and the more fishing sites a bear visited, the more time it spent fishing. Salmon originating from different streams return from sea to spawn at different times, making resource surfing possible.

"Results suggest that in intact watersheds with abundant salmon runs, year to year variation in salmon numbers likely has less effect on [salmon](#) consumption than individual variation in bear foraging behavior," Deacy said. "A remaining challenge is to understand the drivers of that individual variation."

More information: William W. Deacy et al, Phenological tracking associated with increased salmon consumption by brown bears, *Scientific Reports* (2018). [DOI: 10.1038/s41598-018-29425-3](https://doi.org/10.1038/s41598-018-29425-3)

Provided by Oregon State University

Citation: Diverse salmon populations enable 'resource surfing' bears to eat tons of fish (2018, July 23) retrieved 3 May 2024 from <https://phys.org/news/2018-07-diverse-salmon-populations-enable-resource.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--