


# Evidence of *Salmonella Paratyphi C* found for the first time in medieval northern Europe

July 19 2018

---



The face reconstruction of the young woman. Credit: Caroline Wilkinson, Mark Roughley, Ching Liu and Kathryn Smith at the Face Lab at Liverpool John Moores University for their different contributions to the production of the facial depiction

Genome research conducted by the University of Warwick suggests that

enteric fever, a potentially lethal disease more commonly found in hot countries, was present in medieval Europe.


*Salmonella Paratyphi C* causes enteric fever, a life-threatening infection, and has been detected in a 800 year old human skeleton discovered in Trondheim, Norway.

Now scientists are speculating that the evolution of enteric fever could be linked to the domestication of pigs across northern Europe.

The research was conducted by a team of international collaborators led by Professor Mark Achtman of the University's Warwick Medical School and their paper Pan-genome Analysis of Ancient and Modern *Salmonella enterica* Demonstrates Genomic Stability of the Invasive Para C Lineage for Millennia has been published in the journal *Current Biology*.

He and his team analysed bacterial DNA found in the teeth and bones of the skeleton of a young woman who is believed to have migrated to Trondheim from the northernmost areas of Scandinavia or Northwest Russia by her early teens only to die there around the age of 19-24 years.

They reconstructed a genome of *Salmonella Paratyphi C* which causes enteric fever in areas of poor sanitation and lack of clean drinking water. Their discovery indicates that the young Norwegian died of this disease and suggests that these bacteria have long caused enteric fever across northern Europe,



The skeleton plus teeth and bones. Credit: University of Warwick

Prof Achtman said: "*Paratyphi C* is very rare today in Europe and North America except for occasional travellers from South and East Asia or Africa, where the disease is more common. This is the first time that any *Salmonella* have been found in old human remains in Europe, which is surprising because other *Salmonella* are more common today, including *Salmonella* causing typhoid fever, called Typhi, and *Salmonella* causing food poisoning. Earlier this year, Vågene and co-authors described related *Paratyphi C* from skeletons in Mexico, who died in 1545 CE, and speculated the *Paratyphi C* entered the Americas together with Europeans."

The new results included comparative analyses of the *Paratyphi C* genome found in the skeleton against modern *Salmonella* genome sequences from EnteroBase, an online database developed at the University of Warwick and used internationally. This revealed that *Paratyphi C* represents the evolutionary descendants of a common ancestor, or clade, within the Para C lineage. The Para C Lineage includes Choleraesuis, which causes septicaemia in pigs and boar and Typhisuis which causes epidemic swine salmonellosis (chronic paratyphoid) in domestic pigs. These different host specificities likely evolved in Europe over the last 4,000 years and coincide with the timing of pig domestication in Europe.

According to historical records, humans have long been afflicted by bacterial infections, yet genomic analyses of living bacterial pathogens routinely estimate a date for the most recent common ancestor of no more than a few centuries. In general, evolutionary trees contain a stem group, which may include lineages that are now rare or extinct, as well as the crown group of living organisms. Historical reconstructions based only on the crown group ignore the older sub-lineages in the stem group and thereby provide an incomplete picture of the older evolutionary history of the pathogen. In contrast, analyses of ancient DNA such as the *Paratyphi C* genome can shed light on additional millennia of bacterial pathogen evolution that occurred prior to the origin of the crown group.

Professor Achtman added: "Using EnteroBase we were able to define the Para C lineage from 50,000 modern *Salmonella enterica* genomes and find that over its 3,000 year history only a few genomic changes occurred within the Para C lineage.

"As well as reshaping our understanding of *Salmonella enterica*, our research has triggered intriguing speculations about historical host jumps during the Neolithic period between humans and their domesticated animals."

**More information:** Pan-genome Analysis of Ancient and Modern *Salmonella enterica* Demonstrates Genomic Stability of the Invasive Para C Lineage for Millennia, *Current Biology*, [doi.org/10.1016/j.cub.2018.05.058](https://doi.org/10.1016/j.cub.2018.05.058)

Provided by University of Warwick

Citation: Evidence of *Salmonella Paratyphi C* found for the first time in medieval northern Europe (2018, July 19) retrieved 20 April 2024 from <https://phys.org/news/2018-07-evidence-salmonella-paratyphi-medieval-northern.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.