

High climate variability and increasing aridity brought an end to an early hominid species

July 10 2018




The catchment area of the Limpopo River. Here lived an ancestor of man, *Paranthropus robustus* about two million years ago. Credit: MARUM – Center for Marine Environmental Sciences at the University of Bremen

Africa plays a prominent role in human evolution, and is considered by researchers to be the cradle of humanity. In the mid-20th century anthropologists found fossils of *Paranthropus robustus* in South Africa, which belongs to an evolutionary side branch of *Homo sapiens*. *Paranthropus robustus* lived around two million years ago but eventually died out. Possible reasons for the extinction have now been brought to light by an international team of anthropologists and geoscientists led by

Dr. Thibaut Caley of the University of Bordeaux. In this effort, the researchers, who include Dr. Lydie Dupont and Dr. Enno Schefuß of MARUM – Center for Marine Environmental Sciences at the University of Bremen combined various indicators to reconstruct the climatic conditions in southeast Africa at this time. The results were now published in the professional journal *Nature*.

The idea of combining different methods arose from a seeming contradiction: While [climate](#) records from northern Africa indicate that conditions were becoming more arid, data from Lake Malawi suggest exactly the opposite. Was it really true that southeast Africa became more humid as northern Africa became drier? What, then, could have led to the extinction of *Paranthropus robustus*? Lake Malawi lies to the northeast of the catchment basin of the Limpopo, one of Africa's largest rivers. In Maputo Bay (Mozambique) the Limpopo flows into the Indian Ocean. The sediment core – the archive investigated by the researchers for this study – was taken from here.

Due to the continuous deposition, marine sediment cores allow the researchers to observe a sequence of climate changes over long periods of time. Microfossils and pollen from the land are also washed into the ocean by the Limpopo and deposited on the ocean floor. This allows the findings from sites on land to be compared to their temporal development. As Dr. Lydie Dupont from MARUM notes, data from the land often only encompass short time periods, but they can provide important clues about the occurrences of species and their food sources. With the help of the [sediment cores](#), the scientists had access to a climate record spanning about 2.14 million years.



The original complete skull of a 1.8 million years old *Paranthropus robustus* discovered in South Africa. Collection of the Transvaal Museum, Northern Flagship Institute, Pretoria South Africa. Credit: José Braga, Didier Descouens; Ditsong National Museum of Natural History; licensed under CC BY-SA 4.0

The team combined very different kinds of analyses. Hydrogen as well as carbon isotopes of molecular plant fossils were investigated, and these were compared with the results of pollen analyses and element compositions. Each analysis alone can be interpreted in different ways. "Only through the comprehensive consideration could a coherent picture

of the climate in the Limpopo region be reconstructed," says Lydie Dupont.

Furthermore, the team determined the changes in sea-surface temperatures for this time period, allowing them to assess the influence of the ocean on the climate on land. Combining their results with data from the literature, the researchers were able to draw conclusions about the causes of climate changes that occurred during the time when *Paranthropus robustus* lived and ultimately died out.

The combined results from Limpopo portray a different picture than the study from Lake Malawi. From about 1 million years ago to 600,000 years ago climate became substantially more arid. At the same time, climate variability increased noticeably. "What eventually led to the extinction is difficult to say," says Dr. Enno Schefuß. Climatic changes always lead to adaptations by organisms – including adaptation of their feeding habits. If the conditions change extremely rapidly over a short period of time, organisms are less apt to adjust evolutionarily to the altered circumstances. According to the findings in the Limpopo region, *Paranthropus robustus* died out 600,000 years ago.

More information: Thibaut Caley et al. A two-million-year-long hydroclimatic context for hominin evolution in southeastern Africa, *Nature* (2018). [DOI: 10.1038/s41586-018-0309-6](https://doi.org/10.1038/s41586-018-0309-6)

Provided by University of Bremen

Citation: High climate variability and increasing aridity brought an end to an early hominid species (2018, July 10) retrieved 11 May 2024 from <https://phys.org/news/2018-07-evolution-climate-southeast-africa.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.