

# Great Barrier Reef not bouncing back as before, but there is hope

July 18 2018

---



Reefs experiencing flood and poor water quality. Credit: Peter Mumby

The Great Barrier Reef is losing its ability to recover from disturbances, but effective local management could revive its capacity to bounce back.

Scientists at The University of Queensland, ARC Centre of Excellence for Coral Reefs Studies (Coral CoE) and the Australian Institute of Marine Science (AIMS) have found a decline in the ability of Great


Barrier Reef Marine Park reefs to recover after bleaching events, outbreaks of crown of thorns starfish or cyclones over an 18-year period, from 1992 to 2010, even before the recent back-to-back bleaching in 2016 and 2017.

Dr. Juan Ortiz, lead author from The Australian Institute of Marine Sciences and UQ's School of Biological Sciences, said that during this time, average coral [recovery](#) rates showed a six-fold decline across the Great Barrier Reef.

"This is the first time a decline in recovery rate of this magnitude has been identified in coral reefs," he said.

The decline is driven by a combination of the legacy effect of acute disturbances like [coral bleaching](#) and cyclones and the ongoing effect of chronic pressures like poor water quality and [climate change](#).

Professor Peter Mumby of the ARC Centre of Excellence for Coral Reef Studies at The University of Queensland, said that this was serious cause for concern, particularly given the accelerating impacts of climate change on reefs, but it is important to stress that not all reefs are failing.



Reefs experiencing moderate levels of sediment deposition. Credit: Peter Mumby

"I believe there is scope for management to help remedy the situation," he said.

"Our results indicate that coral recovery is sensitive to water quality, and is suppressed for several years following powerful cyclones.

"Some reefs could improve their recovery ability if the quality of the water entering the [reef](#) is actively improved."

Study co-author Dr. Nicholas Wolff, from The Nature Conservancy, said that some areas of the reef are faring better than others, but their overall finding was that action needs to be taken.



Healthy coral reef after recovery. Credit: Peter Mumby

"While there was variability among regions, the decline in recovery rate was consistent in all coral types included in the study," he said.

Dr. Ortiz said that the frequency of acute disturbances was predicted to increase, making careful management key.

"The future of the Great Barrier Reef is threatened without further local management to reduce chronic disturbances and support recovery, and strong global action to limit the effect of climate change."

**More information:** J.-C. Ortiz et al., "Impaired recovery of the Great Barrier Reef under cumulative stress," *Science Advances* (2018). DOI: [10.1126/sciadv.aar6127](https://doi.org/10.1126/sciadv.aar6127) , [advances.sciencemag.org/content/4/7/eaar6127](https://advances.sciencemag.org/content/4/7/eaar6127)

Provided by University of Queensland

Citation: Great Barrier Reef not bouncing back as before, but there is hope (2018, July 18)  
retrieved 25 April 2024 from <https://phys.org/news/2018-07-great-barrier-reef.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.