

Meningococcal bacterial aggregates form a thick honey-like liquid that flows through blood vessels

July 10 2018

The Inserm team led by Guillaume Duménil at the Institut Pasteur, in collaboration with several teams of physicists, has unraveled a key stage in infection by Neisseria meningitidis, a human pathogen responsible for meningitis in infants and young adults. Bacterial aggregates in blood vessels appear to facilitate the progression of the disease. Even if treatment is administered rapidly, the mortality rate due to meningococcal infections remains very high.

Human meningococcal meningitis is characterized by bacterial accumulation in blood vessels, which completely fill up with bacteria. But the mechanisms that govern the formation of these aggregates and the impact of the process are still unknown. A consortium of scientists, intrigued by this formation of intravascular aggregates, set out to understand this stage of infection, especially its underlying physical basis. "The bacterial aggregates formed by Neisseria meningitidis unexpectedly behave like a thick liquid, with a consistency similar to that of honey," explains Guillaume Duménil, head of the Pathogenesis of Vascular Infections Unit. "The bacteria multiply rapidly in blood vessels, forming aggregates which gradually adapt to the complex geometry of the vascular network, like a flowing liquid." The research shows that the formation of these aggregates and their viscous properties are vital for the progression of the infection. A bacterial mutant that forms solid rather than liquid aggregates is at a considerable disadvantage in colonizing blood vessels.



A viscous liquid with original properties

The thick liquid nature of the aggregates depends on a virulence factor known as type IV pili. These long filaments, which have adhesive and dynamic properties, are constantly extending and retracting at the bacterial surface. The bacteria use type IV pili to find other bacteria, draw them in and temporarily establish contact. Aggregation is therefore based on an intermittent process of attraction between bacteria—in other words, a constant alternation between the presence and absence of attraction. In physical terms, this intermittent process of interaction gives the aggregates original properties that had not previously been described. For example, bacteria inside aggregates exhibit a higher level of motility than that observed in the diffusion of isolated bacteria. "As well as improving our understanding of a lethal human infection, our research reveals a new type of active matter—a bacterial aggregate with a viscous, honey-like consistency—based on the intermittent attractive forces between its component parts," concludes Guillaume Duménil.

This multidisciplinary study was the result of close collaboration between a laboratory specializing in meningococcal infections (Guillaume Duménil, Institut Pasteur and Inserm) and physicists. By working with the teams led by Nelly Henry (CNRS, UPMC), Raphael Voituriez (CNRS, UPMC) and Hugues Chaté (CEA, CNRS, Paris-Saclay University), the researchers were able to combine a quantitative experimental approach with a physics-based model of active matter.

More information: Daria Bonazzi et al, Intermittent Pili-Mediated Forces Fluidize Neisseria meningitidis Aggregates Promoting Vascular Colonization, *Cell* (2018). <u>DOI: 10.1016/j.cell.2018.04.010</u>

Provided by Pasteur Institute



Citation: Meningococcal bacterial aggregates form a thick honey-like liquid that flows through blood vessels (2018, July 10) retrieved 7 May 2024 from https://phys.org/news/2018-07-meningococcal-bacterial-aggregates-thick-honey-like.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.