

# Mysteries of Okinawan habu venom decoded

July 26 2018

---



The Okinawan habu is endemic to the Ryukyu islands of Japan. It is the longest and the most venomous of the three habu species found in the region. Credit: Prof. Hiroki Shibata

It is more likely to see a habu snake in a bottle of Okinawan rice liquor

than to see one slithering by the wayside. Even so, habus are very common in the Ryukyu islands, of which Okinawa is a part. Okinawa is home to three species of habu; the Okinawan habu (*Protobothrops flavoviridis*), endemic to the Ryukyu islands, is highly venomous.

A collaborative team of scientists from many research institutes and universities across Japan has now mapped the entire genome of the Okinawan habu, including the genes involved in venom production. Professor Nori Satoh and his team of researchers from Okinawa Institute of Science and Technology Graduate University (OIST), were a part of the study, which was published in *Scientific Reports*. Snake venom has attracted the attention of many scientists worldwide as it has tremendous potential in the development of new drugs. The findings from this study might advance efforts to identify novel therapies for treating ailments such as cancer, cardiovascular diseases and many others.

For the study, the researchers caught Okinawan habus in the wild using a long metal pole with a hook at one end, and collected venom, blood and tissue samples. Then, using the cutting-edge DNA sequencing facilities at OIST, the researchers identified nearly 60 genes from 18 gene families that help the snake make its venom.

Nearly 50 people—most often farm workers tending crops in fields and orchards—are reportedly bitten by habus every year in Okinawa alone. The venom of the Okinawan habu is hemotoxic, destroying blood cells and tissues. Though antidotes are now available, bites can still cause severe pain and permanent tissue damage in humans. The scientists hope the results of this study will contribute to more effective antidotes and treatments for habu bites. "The findings of this study are very important, especially to the people of Okinawa and the surrounding islands where the Okinawan habu is found," says Prof. Satoh.



The Okinawan Habu can produce a large amount of venom. The venom is hemotoxic and destroys the blood cells and tissues of the prey. Credit: Prof. Hiroki Shibata

Snake venom is a cocktail of complex proteins. "Though the Okinawan habu is not as venomous as cobras or taipans, it can produce a large amount of venom—up to 1 ml," says Prof. Hiroki Shibata, a collaborator from the University of Kyushu. Its long, sharp fangs help discharge the venom deep inside the prey, he explains.

The genome of any organism is a treasure trove of information. Mapping venom genes of the habu not only allows scientists to explore its potential in drug development, but also gives scientists the tools to study how venom toxins have evolved, their composition, and the mechanism by which they do their damage.

The researchers explored this venom's evolutionary history by constructing a phylogenetic tree of venom genes, including those that they identified in the habu. A phylogenetic tree helps illustrate the relationship between two or more organism groups by identifying the similarities and dissimilarities between them. The distance between the groups in a tree translates to different degrees of relatedness between them. The groups that are closely placed are more related to one another than those placed further away.

Venomous snakes around the world belong to one of the two families; Viperidae or Elapidae. The members of the family Viperidae, including the habu, have hemotoxic venom, whereas the members of the family Elapidae, such as cobras, have venom that is neurotoxic. In this study, the researchers found many common components among venom genes of both families, thereby tracing back the origin of the present-day venom toxins to a common ancestor that existed millions of years ago. "The venom gene copies originated a long time ago, possibly at the early stage of vertebrate evolution," explains Prof. Shibata. The scientists believe that combined information gathered from venom research so far will help unravel the enigma behind the process of [venom](#) production in snakes.

Provided by Okinawa Institute of Science and Technology

Citation: Mysteries of Okinawan habu venom decoded (2018, July 26) retrieved 27 January 2023 from <https://phys.org/news/2018-07-mysteries-okinawan-habu-venom-decoded.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.