

Pieces of mantle found rising under north and south ends of Cascadia fault

July 25 2018



Map shows the Cascadia Subduction Zone along the Pacific Northwest coast, with a shaded area encompassing the onshore and offshore areas where seismometers were located. Data from the seismometers helped University of Oregon researchers identify seismic anomalies at both ends of the fault where they believe pieces of the upper mantle are rising and modulating earthquake activity. Credit: Miles Brodmer, University of Oregon

With four years of data from 268 seismometers on the ocean floor and



several hundred on land, researchers have found anomalies in the upper mantle below both ends of the Cascadia Subduction Zone. They may influence the location, frequency and strength of earthquake events along the U.S. Pacific Northwest.

The anomalies, which reflect regions with lower <u>seismic wave velocities</u> than elsewhere beneath the fault line, point to pieces of the Earth's upper mantle that are rising and buoyant because of melting rock and possibly elevated temperatures, said Miles Bodmer, a University of Oregon doctoral student who led a study now online as an accepted paper by the journal *Geophysical Research Letters*.

The 620-mile subduction zone, which hasn't experienced a massive lengthwise <u>earthquake</u> since 1700, is where the Juan de Fuca ocean plate dips under the North American continental plate. The fault zone stretches just offshore from northern Vancouver Island to Cape Mendocino in northern California.

The mantle is rising under the southern Gorda deformation zone at the north edge of the San Andreas Fault and under the Olympic Peninsula and southern Vancouver Island.

"What we see are these two anomalies that are beneath the subducting slab in the northern and southern parts of the subduction zone," Bodmer said. "These regions don't have the same behavior as the entire fault. There are three segments that have their own distinct geological characteristics. The north and south segments have increased locking and increased tremor densities."

Locking refers to how strongly two plates stick. "If they are stuck together tightly, as is the case here, they are building up stress, and you have the potential for the release of that stress, or energy, in large earthquake events," Bodmer said.



Such quakes, while strong, are below that of the 9-plus magnitude event projected should all of Cascadia rupture at once, he said. Locking is much weaker in Cascadia's central section, which includes most of Oregon, where infrequent, smaller quakes tend to occur from creeping along the plates.

Tremor refers to long-duration seismic signals often seen at subduction zones. "These happen deep and take more time than a typical earthquake as they rumble to release energy," Bodmer said.

The findings won't help earthquake forecasting, but they do point to the need for real time onshore-offshore seismic monitoring and geodetic analyses, such as from GPS to help plot spatial coordinates, of the anomalies as a next step in that direction, said co-author Douglas R. Toomey, a seismologist in the UO Department of Earth Sciences.

The study helps to make sense of Cascadia's historical record of earthquake, he said.

The junction of the Cascadia-San Andreas faults, Toomey said, contains a lot of complexity and is the most seismically active part of contiguous North America. Seismic history also shows more earthquake activity in the Puget Sound area than in central Oregon. Both regions accumulate energy that eventually is released in large earthquakes, he said.

"Our study is worse news for Portland northward to Seattle and for southern Cascadia, but central Cascadia is not off the hook," said Toomey, who also is lead investigator for the Oregon component of ShakeAlert, the West Coast early warning network. "More frequent earthquakes to the north and south are seen in historical seismicity patterns. This research helps to understand that."

The study involved deep imaging, similar to CAT scans, using different



forms of seismic waves coming from distant earthquakes moving through the Earth.

The ocean-bottom seismic stations, from which data were retrieved every 10 months, were part of the National Science Foundation-funded Cascadia Initiative. Older data from numerous onshore studies in the western United States also were included in the analysis.

In addition to helping to understand Cascadia's historical earthquake record, the anomalies, Bodmer said, suggest that the two buoyant ends help to modulate plate coupling forces.

"We're looking at structures deep within the Earth and finding evidence suggesting that they are influencing the megathrust faults and controlling where we see increases in locking and segmentation," Bodmer said. "Knowing the timing and path of the seismic signals, we can look at velocity variation and equate that to the structures. With large offshore data sources, we might be able to better understand how a large rupture in the south might extend into central Oregon."

More information: Miles Bodmer et al, Buoyant Asthenosphere Beneath Cascadia Influences Megathrust Segmentation, *Geophysical Research Letters* (2018). DOI: 10.1029/2018GL078700

Provided by University of Oregon

Citation: Pieces of mantle found rising under north and south ends of Cascadia fault (2018, July 25) retrieved 7 May 2024 from <u>https://phys.org/news/2018-07-pieces-mantle-north-south-cascadia.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private



study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.