


Putting a quantum gas through its phases

July 3 2018



Phase diagrams showing the four different regions observed in the experiment: white -- superfluid without photons; red and yellow -- photons in only one of the cavities; blue -- photons in both cavities simultaneously (mixed phase). As the coupling between the orders is increased, the mixed-phase regime (blue) becomes increasingly favourable. Credit: Esslinger group, ETH Zurich (adapted from doi: 10.1038/s41563-018-0118-1)

Physicists at ETH Zurich have developed an experimental platform for studying the complex phases of a quantum gas characterized by two order parameters. With unprecedented control over the underlying microscopic interactions, the approach should lead to novel insight into the properties of a broad range of fundamentally and technologically important materials.

As a physical system undergoes a phase transition, it typically becomes more or less ordered. For instance, when a piece of iron is heated to above the Curie temperature, the strong ferromagnetic alignment of the elementary magnetic dipole moments gives way to much weaker paramagnetic alignment. Such changes are well described in the general framework of order parameters, provided by the Landau theory of phase transitions. However, many materials of current fundamental and technological interest are characterised by more than one order parameter. And here, the situation can become extraordinarily complex, in particular when the different orders interact with one another. The traditional route to gaining an understanding of such complex quantum systems is to carefully explore the response to changes in external conditions and to various probes, and thus to map out the phase diagram of the system. A complementary approach is now presented by Tobias Donner and his team in the group of Tilman Esslinger in the Department of Physics of ETH Zurich. Their approach controls all relevant microscopic parameters of a quantum system governed by two coupled order parameters, and therefore can essentially construct and modify the [phase diagram](#) from the bottom up, as they report in a paper published today in the journal *Nature Materials*.

Phenomenological models that reproduce the experimentally determined phase diagrams of materials with one or more ordering tendencies have provided deep insight into the behaviour of a variety of systems, such as multiferroics, material that exhibits ferromagnetism and ferroelectric properties, opening the door to new functionality—or certain families of

superconductors. However, the microscopic processes underlying the formation of macroscopic order in these systems remain unknown. This gap in understanding limits the predictive power of phenomenological models, and at the same time, makes it difficult to know just how a given material should be modified to obtain desired properties. Hence the appeal of the approach taken by Donner and his colleagues, who started not with a specific system and its phenomenological description, but with a flexible quantum system whose relevant microscopic parameters can be controlled with high accuracy, and be tuned across a broad range of values, enabling the realization of diverse scenarios.

To create such a versatile platform, the team optically trapped a Bose-Einstein condensate (BEC) at the intersection of two optical cavity modes (see the figure). In this configuration, the BEC can crystallise in two different patterns, each of which is associated with a different order parameter. Depending on the experimental setting, the two orders either competed with one another—forcing the system into one of the two patterns (red and yellow)—or to coexist, leading to a new coupled phase (blue), where the two orders do not simply add, but give rise to a more complex spatial arrangement. The extent of this mixed-order [phase](#) can also be controlled to favour regimes of mutual exclusion or of mutual enhancement.

Whereas these particular phases have no known direct role in practical materials, the approach established with these experiments can be modified in the future to simulate properties of [materials](#) that are technologically relevant. In particular, in cuprate high-temperature superconductors, coupled spin and charge order are known to have an important, yet not fully understood role. The sort of experiments now pioneered by the ETH physicists should offer a unique tool to explore such phases starting from a 'clean' quantum system with well-controlled and widely tunable interactions.

More information: Andrea Morales et al, Coupling two order parameters in a quantum gas, *Nature Materials* (2018). [DOI: 10.1038/s41563-018-0118-1](https://doi.org/10.1038/s41563-018-0118-1)

Provided by ETH Zurich

Citation: Putting a quantum gas through its phases (2018, July 3) retrieved 8 April 2024 from <https://phys.org/news/2018-07-quantum-gas-phases.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
